

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
16 May 2002 (16.05.2002)

(10) International Publication Number  
PCT WO 02/038793 A3

(51) International Patent Classification<sup>7</sup>: C12Q 1/68,  
G01N 33/53, C12N 5/00, C07H 21/04

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,  
ZA, ZW.

(21) International Application Number: PCT/US01/46350

(22) International Filing Date:

2 November 2001 (02.11.2001)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/245,866

7 November 2000 (07.11.2000) US

Published:

— with international search report

(71) Applicant (for all designated States except US): ANADYS  
PHARMACEUTICALS, INC. [US/US]; 9050 Camino  
Santa Fe, San Diego, CA 92121 (US).

(88) Date of publication of the international search report:  
15 August 2002

(72) Inventor; and

(75) Inventor/Applicant (for US only): BICHKO, Vadim  
[US/US]; 23 Clarendon Road, Belmont, MA 02478 (US).

(48) Date of publication of this corrected version:

13 February 2003

(74) Agents: FEHLNER, Paul, F. et al.; Darby & Darby P.C.,  
805 Third Avenue, New York, NY 10022-7513 (US).

(15) Information about Correction:

see PCT Gazette No. 07/2003 of 13 February 2003, Sec-  
tion II

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 02/038793 A3

(54) Title: HEPATITIS C VIRUS CONSTRUCTS CHARACTERIZED BY HIGH EFFICIENCY REPLICATION

(57) Abstract: The present invention relates to the recombinant hepatitis C virus (HCV)-derived nucleic acids and to stable rapidly growing cell clones derived from human hepatoma Huh-7 cell line and supporting high titer replication of said recombinant HCV nucleic acids. The subgenomic HCV replicons and cell clones of the instant invention represent the *in vitro* system of choice for studies of HCV propagation, anti-viral drug screening, and vaccine development.

**HEPATITIS C VIRUS CONSTRUCTS CHARACTERIZED BY**  
**HIGH EFFICIENCY REPLICATION**

**Field of the Invention**

5           The present invention relates to recombinant hepatitis C virus (HCV)-derived nucleic acids and to stable rapidly growing cell clones supporting their efficient replication.

**BACKGROUND OF THE INVENTION**

10           Infection by hepatitis C virus (HCV) is a compelling human medical problem. HCV is recognized as the causative agent for most cases of non-A and non-B hepatitis, with an estimated worldwide prevalence of 170 million cases (*i.e.*, 2-3%) (Choo *et al.*, Science, 1989, 244:359-362; Kuo *et al.*, Science, 1989, 244:362-364; Purcell, FEMS Microbiol. Rev., 1994, 14:181-192; Van der Poel, In: *Current Studies in Hematology and Blood Transfusion*, Reesink ed., Basel: Karger, pp. 137-163, 1994). Four million individuals may be infected in the United States alone (Alter and Mast, Gastroenterol. Clin. North Am., 1994, 23:437-455).

15           HCV is primarily transmitted parenterally, although sexual and perinatal transmission do appear to occur. At present, no risk factor has been identified in about 40% of HCV-infected individuals in the US (Alter, Infect. Agents Dis., 1993, 2:155-166). Upon first exposure to HCV, only about 10% or less of infected individuals develop acute clinical hepatitis, while others appear to resolve the infection spontaneously. In most instances,  
20           however, the virus establishes a chronic infection that persists for decades, leading in about 50% of all cases to chronic hepatitis, which can, in turn, develop into liver cirrhosis and/or

- 5 hepatocellular carcinoma (Iwarson, FEMS Microbiol. Rev., 1994, 14:201-204; Kew, *ibid.* pp.211-220; Saito *et al.*, Proc. Natl. Acad. Sci. USA, 1990, 87:6547-6549).

Apart from liver cells, HCV can also replicate in peripheral blood mononuclear cells (PBMCs) both *in vivo* and in experimentally infected B- and T-cell lines (U.S. Patent Nos.: 5,679,342 and 5,968,775). Such a lymphotropism may account for the numerous  
10 immunological disorders, in particular type II and type III cryoglobulinaemia, observed in more than 50% of chronic hepatitis C patients (Esteban *et al.*, In: *Hepatitis C Virus*, Reesink ed., Basel: Karge, 1998, pp. 102-118).

### HCV Structure and Genome Organization

Given the high prevalence of the virus, HCV has become a focus of intensive research (for recent review *see* Bartenschlager and Lohmann, J. Gen. Virol., 2000, 81:1631-1648;  
15 Rosen and Gretch, Mol. Medicine Today, 1999, 5: 393-399). Originally cloned in 1989 (Choo *et al.*, *supra*), the viral genome is now well characterized. HCV is a (+) strand enveloped RNA virus, *i.e.* its genome is represented by a coding single stranded RNA (cRNA) which is packaged with the structural proteins in a viral particle surrounded by a host cell-derived membrane. HCV has been classified as the sole member of a distinct genus  
20 called hepacivirus in the family *Flaviviridae*, which includes, *e.g.*, the flaviviruses and the animal pathogenic pestiviruses. Its genome has a length of approximately 9.6 kb and its single, long open reading frame (ORF) encodes an approximately 3000-amino acid polyprotein that is proteolytically cleaved into a set of distinct products (Figure 1 [prior art]; *see also* Rice, In: *Virology*, Fields *et al.* eds., Lippincott- Raven, 1996, Vol.1, pp.931-960;  
25 Clarke, J. Gen. Virol., 1997, 78:2397).

The HCV ORF is flanked at the 5' and 3' ends by nontranslated regions (NTRs). Translation of the ORF is directed via an approximately 340 nucleotide (nt) long 5' NTR functioning as an internal ribosome entry site (IRES) and permitting the direct binding of ribosomes in close proximity to the start codon of the ORF (Tsukiyama-Kohara *et al.*, J.  
30 Virol., 1992, 66:1476-1483; Wang *et al.*, J. Virol., 1993, 67:3338-3344). The first approximately 40 nucleotides of the 5' NTR are not required for translation but, based on

5 analogy with other (+) strand RNA viruses, are involved most likely in RNA replication (Boyer and Haenni, *Virology*, 1994, 198:415-426). The 3' NTR has a tripartite structure composed of a variable sequence following the stop codon of the ORF, a poly(U) tract of heterogeneous length and a highly conserved 98 nucleotide sequence essential for replication *in vivo* (Kolykhalov *et al.*, *J. Virol.*, 1996, 70:3363; Tanaka *et al.*, *Biochem. Biophys. Res.*  
10 *Comm.*, 1995, 215:744; Tanaka *et al.*, *J. Virol.*, 1996, 70:3307; Yamada *et al.*, *Virology*, 1996, 223:255; Yanagi *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1999, 96:2291; Kolykhalov *et al.*, *J. Virol.*, 2000, 74:2046-2051).

The HCV polyprotein is cleaved co- and post-translationally by cellular and viral proteinases into ten different products, with the structural proteins located in the N-terminal  
15 one-third and the non-structural (NS) proteins (*i.e.*, proteins which are not expected to be constituents of the virus particle) in the remainder (Figure 1; reviewed in Bartenschlager and Lohmann, *supra*; Bartenschlager, *J. Viral Hepatitis*, 1999, 6:165-181; Reed and Rice, In: *Hepatitis C Virus*, Reesink ed., Basel: Karger, 1998, pp. 1-37). The first cleavage product of the polyprotein is a highly basic core protein, which is the major constituent of the  
20 nucleocapsid (Yasui *et al.*, *J. Virol.*, 1998, 72:6048-605) and is involved in modulation of several cellular processes leading to induction of hepatocellular carcinoma (Chang *et al.*, *J. Virol.*, 1998, 72: 3060-3065; Chen *et al.*, *J. Virol.*, 1997, 71:9417-9426; Matsumoto *et al.*, *ibid.*, pp.1301-1309; Moriya *et al.*, *Nature Med.*, 1998, 4:1065-1067). Envelope proteins E1 and E2 are highly glycosylated type 1 transmembrane proteins, forming two types of stable  
25 heterodimeric complexes (Deleersnyder *et al.*, *J. Virol.*, 1997, 71:697-704). In addition, E2 was shown to interact with the interferon (IFN)-induced double-stranded RNA-activated protein kinase PKR, allowing continuation of translation of HCV RNA in the presence of IFN (Taylor *et al.*, *Science*, 1999, 285:107-110). Protein p7, located at the C-terminus of E2, is a highly hydrophobic polypeptide of unknown function. Most of the nonstructural proteins  
30 NS2-5B are required for replication of the viral RNA (Lohmann *et al.*, *Science*, 1999, 285:110-113). NS2 and the N-terminal domain of NS3 constitute the NS2-3 proteinase, catalysing cleavage at the NS2/3 site (Grakoui *et al.*, 1993, *Proc. Natl. Acad. Sci. USA*, 1993, 90:10583-10587; Hijikata *et al.*, *J. Virol.*, 1993, 67:4665-4675; Hirowatari *et al.*, *Arch. Virol.*, 1993, 133:349-356). NS3 is a bifunctional molecule carrying, in the N-terminal

5 approximately 180 residues, a serine-type proteinase responsible for cleavage at the NS3/4A, NS4A/B, NS4B/5A and NS5A/B sites and, in the C-terminal remainder, NTPase/helicase activities essential for translation and replication of the HCV genome (Bartenschlager *et al.*, J. Virol., 1993, 67:3835-3844; Eckart *et al.*, Biochem. Biophys. Res. Comm., 1993, 192:399-406; Grakoui *et al.*, J. Virol., 1993, 67:2832-2843; Gwack *et al.*, Biochem. Biophys. Res. Comm., 1996, 225:654-659; Hong *et al.*, J. Virol., 1996, 70:4261- 4268; Kim *et al.*, Biochem. Biophys. Res. Comm., 1995, 215:160-166; Suzich *et al.*, J. Virol., 1993, 67:6152-6158; Tai *et al.*, J. Virol., 1996, 70:8477-8484; Tomei *et al.*, J. Gen. Virol., 1993, 77:1065-1070; Kolykhalov *et al.*, 2000, *supra*). In addition, NS3 may interfere with host cell functions by inhibiting protein kinase A (PKA)-mediated signal transduction and/or by  
15 inducing cell transformation (Borowski *et al.*, Eur. J. Biochem., 1996, 237:611-618; Sakamuro *et al.*, J. Virol., 1995, 69:3893-3896). NS4A is an essential cofactor of the NS3 proteinase and is required for efficient polyprotein processing (Bartenschlager *et al.*, J. Virol., 1994, 68:5045- 5055; Failla *et al.*, *ibid.*, pp. 3753-3760; Lin *et al.*, *ibid.*, pp. 8147-8157; Tanji *et al.*, J. Virol., 1995, 69:1575- 1581). The function of the hydrophobic NS4B is so far  
20 unknown. NS5A is a highly phosphorylated protein (Asabe *et al.*, J. Virol., 1997, 71:790-796; Kaneko *et al.*, Biochem. Biophys. Res. Comm., 1994, 205:320-326; Koch and Bartenschlager, J. Virol., 1999, 73:7138- 7146; Neddermann *et al.*, *ibid.*, pp. 9984-9991; Tanji *et al.*, J. Virol., 1995, 69:3980- 3986) which appears to interfere with the antiviral effect of IFN by binding to PKR (Gale *et al.*, Virology, 1997, 230:217-227; Gale *et al.*, Mol. Cell  
25 Biol., 1998, 18:5208-5218) and may also play role in RNA replication. NS5B was identified as the RNA-dependent RNA polymerase (RdRp) (Al *et al.*, Virus Res., 1998, 53:141-149; Behrens *et al.*, EMBO J., 1996, 15:12-22; Lohmann *et al.*, J. Virol., 1997, 71:8416-8428; Yamashita *et al.*, J. Biol. Chem., 1998, 273:15479-15486; Yuan *et al.*, Biochem. Biophys. Res. Comm., 1997, 232:231-235).

### HCV Replication and Variability

30 Similarly to related positive (+) strand RNA viruses, HCV replication occurs by means of a negative (-) strand RNA intermediate and is catalyzed by the NS proteins forming

5 a cytoplasmic membrane-associated replicase complex. HCV replication cycle can be summarized as follows (Figure 2 [prior art]): (1) penetration of the host cell and liberation of the genomic RNA (cRNA) from the virus particle into the cytoplasm; (2) translation of the input cRNA, processing of the polyprotein and formation of a replicase complex associated with intracellular membranes; (3) utilization of the input (+) strand for synthesis of a (-)  
10 strand RNA intermediate; (4) production of new (+) strand RNA molecules which can be used for synthesis of new (-) strands, for polyprotein expression or for packaging into progeny virions; (5) release of virus from the infected cell via cellular secretion pathway resulting in formation of cell-derived viral membrane envelope.

The dynamics of HCV replication can be deduced from the rapid rates of virus  
15 production and emergence of mutants. Analysis of viral dynamics during antiviral treatment of patients with IFN- $\alpha$  revealed a virion half-life of 3–5 hours and a clearance and production rate of approximately  $10^{12}$  particles per day (Zeuzem *et al.*, Hepatology, 1998, 28:245-252; Neumann *et al.*, Science, 1998, 282:103-107; Ramratnam *et al.*, Lancet, 1999, 354:1782-1785). Although in absolute amounts these numbers are high, they are not with  
20 respect to a single cell, corresponding to a virion production rate of 50 particles per hepatocyte per day (Neumann *et al.*, *supra*).

Another feature of HCV replication is a rapid generation of virus variants. Early studies of the mutation rate of HCV in chronically infected humans and chimpanzees demonstrated that this virus mutated very rapidly with the rate of change varying between  
25 different genomic regions (Ogata *et al.*, Proc. Natl. Acad. Sci. USA, 1991, 88:3392-3396; Okamoto *et al.*, Virology, 1992, 190:894-899). Thus, the highest mutation rate was found in the E1 and E2 genes with especially high rate observed in a short sequence encoding the domain located at the N-terminus of E2 protein (hence termed “hypervariable region 1” or “HVR1”). Accordingly, even within a single patient HCV does not exist as a single entity  
30 but rather as a collection of microvariants of a predominant “master sequence”, a phenomenon that has been referred to as *quasispecies* (reviewed in Bukh *et al.*, Semin. Liv. Dis., 1995, 15:41-63; Bukh *et al.*, Clin. Exp. Rheumatol., 1995, 13(suppl.):S3-S7; Holland *et al.*, Curr. Topics Microbiol. Immunol., 176:1- 20). The master sequence, as well as the

5 consensus sequence of the quasispecies sequence population have been found to change sequentially during the infection. The production of such large number of variants is primarily due to the high error rate of the viral RdRp that, based on analogies with RdRps of other (+) strand RNA viruses, is expected to be in the range of  $10^{-4}$ . Using comparative  
10 sequence analyses of HCV genomes isolated over intervals of 8 or 13 years, a mutation rate of  $1.44 \times 10^{-3}$  or  $1.92 \times 10^{-3}$  base substitutions per site per year was found, respectively (Ogata *et al.*, *supra*; Okamoto *et al.*, *supra*). The high variation observed with HCV replication may also account for the fact that a significant fraction of virus genomes appear to be defective (Martell *et al.*, J. Virol., 1992, 66:3225-3229).

The biological consequences of quasispecies include: (i) the development of escape  
15 mutants to humoral and cellular immunity leading to the establishment of a persistent infection; (ii) variable cell tropism (*e.g.*, lymphotropic vs hepatotropic); (iii) vaccine failure, and (iv) rapid development of drug resistance (Bukh *et al.*, Semin. Liv. Dis., *supra*). For example, it has been found that the HVR1 contains epitopes that elicit a specific humoral immune response and that sequential changes of HVR1 during infection resulted in the  
20 emergence of epitopes that were not recognized by pre-existing antibodies (Weiner *et al.*, Proc. Natl. Acad. Sci. USA, 1992, 89:3468-72; Taniguchi *et al.*, Virology, 1993, 195:297-301; Kato *et al.*, J. Virol., 1993, 67:3923-30; Kato *et al.*, J. Virol., 1994, 68:4776-84).

It is now well established that HCV exists as distinct genotypes among different HCV isolates with prevalence of each of the genotypes in specific geographical locations. Based  
25 on the genomic variability in the most highly conserved NS5B and E1 sequences, HCV has been classified into at least 9 major genetic groups (genotypes 1a, 1b, 1c, 2a, 2b, 3a, 7, 8, 9) with total over 30 subtypes (Bukh *et al.*, Clin. Exp. Rheumatol., *supra*; Simmonds *et al.*, J. Gen. Virol., 1993, 74:2391-2399). Several recent studies indicate that the extensive genetic heterogeneity of HCV may have important clinical implications, with genotype 1b (prevalent  
30 in the US and Europe) being associated with a more severe liver disease and a poorer response to interferon therapy (reviewed in Bukh *et al.*, *supra*).

### Anti-HCV Therapies

5        Despite the intense research, the only anti-HCV therapy currently available is administration of a high dose of IFN- $\alpha$  or a combination treatment with IFN- $\alpha$  and the nucleoside analogue ribavirin. However, only about 40% of all patients benefit from this treatment and develop a sustained response, demonstrating the urgent need for more effective antiviral therapeutics (Marcellin *et al.*, *Ann. Intern. Med.*, 1997, 127:875-881; Reichard *et al.*,  
10    *Lancet*, 1998, 351:83-87).

      As mentioned above, anti-HCV vaccine development has been hampered by the high degree of viral variability leading to efficient immune evasion and the lack of protection against reinfection, even with the same inoculum (Farci *et al.*, *Science*, 1992, 258:135-140; Kao *et al.*, *J. Med. Virol.*, 1996, 50:303-308; Shimizu *et al.*, 1994, *J. Virol.*, 68:1494-1500;  
15    Wyatt *et al.*, *J. Virol.*, 1998, 72:1725-1730). Among the most promising approaches to vaccine development, is immunization with highly conserved HCV core protein alone or in combination with E1 and/or E2 envelope proteins, and/or NS3 protein (Major *et al.*, 1995, *J. Virol.*, 69:5798-5805; Tokushige *et al.*, *Hepatology*, 1996, 24:14-20; Geissler *et al.*, *J. Immunol.*, 1997, 158:1231-1237; Inchauspe *et al.*, *Vaccine*, 1997, 15:853-856).

20        In view of the reasonable scepticism over the "universal" anti-HCV vaccine, the importance of alternative therapies increases. Such therapies include without limitation: (i) small molecule inhibitors directed against specific viral targets (*e.g.*, E1/E2 envelope proteins or NS3 protease/helicase); (ii) antisense oligonucleotides and ribozymes for the inhibition of HCV replication, and (iii) transdominant-negative proteins (Kim *et al.*, *Cell*, 1996, 87:343-  
25    355; Love *et al.*, *ibid.*, 331-342; Yao *et al.*, *Nat. Struct. Biol.*, 1997, 4:463-467; Yan *et al.*, *Protein Sci.*, 1998, 7:837-847; Von Wilsaker *et al.*, *Hepatology*, 1997, 26:251-255; Lieber *et al.*, *J. Virol.*, 1996, 70:8782-8791; Rosen and Gretch, *supra*; Saito *et al.*, *Gastroenterology*, 1997, 112:1321-1330; Nakano *et al.*, *J. Virol.*, 1997, 71:7101-7109; Fournillier *et al.*, *Hepatology*, 1998, 28:237-244; Wakita *et al.*, *J. Biol. Chem.*, 1994, 269:14205-14210;  
30    Mizutani *et al.*, *Biochem. Biophys. Res. Comm.*, 1995, 212:906-911; Alt *et al.*, *Hepatology*, 1995, 22:707-717; Havecak *et al.*, *J. Virol.*, 1996, 70:5203-5212; Lima *et al.*, *J. Biol. Chem.*, 1997, 272:626-638; Alt *et al.*, *Arch. Virol.*, 1997, 142:589-599; Wu and Wu,



- 5 Gastroenterology, 1998, 114:1304-1312; Sakamoto *et al.*, J. Clin. Invest., 1996, 98:2720-2728; Ruiz *et al.*, J. Viral Hepatitis, 1999, 6:17-34).

### *In Vivo and In Vitro Systems for HCV Infection*

As the currently available chimpanzee models are too expensive to be practical for early stage evaluation of anti-HCV therapeutics, there is a need in the art to develop more manageable and efficient *in vivo* and *in vitro* model systems for HCV infection.

- 10 In an attempt to establish a small animal model Xie *et al.* (Virology, 1998, 244:513-520) experimentally inoculated *Tupaia*s (*T. belangeri chinensis*), a tree shrew species which adapt and breed in the laboratory environment, are closely related to primates, and were previously shown to be susceptible to infection with the human rotavirus (Wan *et al.*, Natl. Med. J. Chin., 1982, 62:461-465), herpes simplex virus type 1 and 2 (Darai *et al.*, J. Infect. Dis., 1978, 137:221-226), and human hepatitis viruses A, B (both *in vitro* and *in vivo*), and Delta (Li *et al.*, Chung. Hua. I. Hsueh. Tsa. Chih., 1995, 75:611-613; Walter *et al.*, Hepatology, 1996, 24:1-5; Yan *et al.*, J.Cancer Res. Clin. Oncol., 1996, 122:283-288 and 289-295; Zan *et al.*, Acta Acad. Med. Sin., 1981, 3:148-152). Although, upon inoculation, only about one-quarter of the animals became infected with HCV and developed either  
20 transient or intermittent viraemia with rather low titers, the *Tupaia* animal model appears to be very promising. Recently, two potential alternatives have been described for the propagation of hepatitis B viruses that might be used for HCV as well. Both systems are based on the engraftment of human liver tissue into immuno-compromised mice (Ilan *et al.*, Hepatology, 1999, 29:553-562; Petersen *et al.*, Proc. Natl. Acad. Sci. USA, 1998, 95:310-315).  
25

- Although development of a small animal model is critical for studies of HCV pathophysiology and for assaying toxicology and pharmacokinetics of anti-HCV therapeutics, it is of primary importance to create a convenient and reliable cell culture-based assay system that supports HCV infection and replication and allows detailed molecular studies of HCV  
30 propagation and efficient high-throughput evaluation of anti-HCV therapeutics.

5           Until recently, *in vitro* research on HCV has depended largely on (i) analogies to the closely related flavi- and pestiviruses, (ii) characterization of recombinantly produced HCV proteins, (iii) infection of primary cell cultures with HCV-containing sera of infected individuals, and (iv) cultivation of primary cells derived from chronically infected tissues (Lanford *et al.*, 1994, *Virology*, 202:606; Shimizu *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1992, 10   89:5477; Mizutani *et al.*, *J. Virol.*, 1996, 70:7219-7223; Ikeda *et al.*, *Virus Res.*, 1998, 56:157; Fournier *et al.*, *J. Gen. Virol.*, 1998, 79:2376). For example, HCV replication was reported in primary hepatocytes from humans or chimpanzees following infection with high titer HCV- containing serum (Fournier *et al.*, *supra*; Iacovacci *et al.*, *Res. Virol.*, 1993, 144:275-279; Lanford *et al.*, *supra*; Rumin *et al.*, *J. Gen. Virol.*, 1999, 80:3007-3018) and in 15   PBMCs isolated from chronically infected patients (Bouffard *et al.*, *J. Infect. Diseases*, 1992, 166:1276-1280; Müller *et al.*, *J. Gen. Virol.*, 1993, 74:669-676; Zignego *et al.*, *J. Hepatol.*, 1992, 15:382-386).

          In the primary cell cultures HCV replication was detected using a number of highly sensitive methods: (i) detection of (-) strand replicative RNA intermediates by strand-specific 20   reverse transcriptase-polymerase chain reaction (RT-PCR) or Northern blot; (ii) determination of an increase of (+) strand RNA during the cultivation period using either b-DNA assay or quantitative RT-PCR; (iii) detection of inhibition of replication upon incubation of the cells with IFN- $\alpha$  or antisense oligonucleotides; (iv) the ability of the cell culture medium (containing viral particles) to cause infection upon addition to naive cells; (v) sequence 25   analysis of HCV genomes or genome fragments to demonstrate genomic variability and selection of variants upon infection and cultivation, and (vi) detection of viral antigens *in situ* or upon isolation using, *e.g.*, immunofluorescence, Western blotting, or flow cytometry.

          It should be noted, however, that primary cell cultures are not routinely available, are hard to maintain, and suffer from poor reproducibility and a low level of HCV replication that 30   can be measured only with highly sensitive techniques. Accordingly, numerous attempts have been made to develop stable HCV-replicating cell lines. Initially, these cell lines were also developed by infecting them with the virus isolated from infected individuals. Such HCV-infected cell lines, secondary monkey kidney cells CV-1 and human diploid fibroblasts

5 VERO, are disclosed, for example, in the PCT Application No. WO 96/24662. With respect to hepatoma cell lines, the most detailed results are available for the non-neoplastic cell line PH5CH (Ikeda *et al.*, 1998, *supra*; Kato *et al.*, Jap. J. Cancer Res., 1996, 87:787-792). However, a strong selection for HCV variants in the HVR1 of the E2 protein was observed in this cell line suggesting that only certain variants can bind to or replicate in these cells. A  
10 similar decrease of complexity of viral quasispecies has been described for HCV propagated in primary human hepatocytes for up to 3 months (Rumin *et al.*, *supra*).

HCV replication upon infection was also demonstrated in cultured cells derived from the T-cell lines MT-2, HPBMa10-2, and MOLT-4 and from the B-cell line Daudi. For example, in HPBMa10-2 and Daudi cells, long-term propagation of HCV for more than 1  
15 year has been described (Nakajima *et al.*, J. Virol., 1996, 70:3325-3329), and virus could be transmitted several times to naive cells by cocultivation (Shimizu and Yoshikura, J. Virol., 1994, 68:8406-8408). HCV-replicating B- and T-cell lines are also disclosed in the U.S. Patent No. 5,679,342. However, similarly to PH5CH cells, it was found that only certain virus variants replicate in HPBMa10-2 and Daudi cells as well as in MT-2C cells, suggesting  
20 the selection of lymphotropic HCV quasispecies (Mizutani *et al.*, J. Virol., 1996, *supra*; Sugiyama *et al.*, J. Gen. Virol., 1997, 78:329-336).

#### **Attempts to Create Stable Cell Lines Containing Self- Propagating Recombinant HCV Replicons**

The recent construction of cloned HCV genomes and demonstration of their ability to replicate and cause disease development after intrahepatic inoculation of chimpanzees (Beard  
*et al.*, Hepatology, 1999, 30:316- 324; Kolykhalov *et al.*, Science, 1997, 277:570-574;  
25 Yanagi *et al.*, Proc. Natl. Acad. Sci. USA, 1997, 94:8738; Yanagi *et al.*, Virology, 1998, 244:161) has opened some new avenues to study HCV replication and pathogenesis. Specifically, it made feasible the development of stable cell cultures containing selectable HCV replicons. Compared to the infection of cell lines with HCV- containing patient material, the introduction of cloned virus genomes is superior because the inoculum is well  
30 defined and can be generated in high quantities. Most importantly, the genome can be

5 manipulated at will, permitting a detailed genetic analysis of viral functions leading to successful development of anti-viral therapeutics.

Similarly to experience with several (+) strand RNA viruses (Boyer and Haenni, *supra*), it became clear with the first attempts to create self-replicating subgenomic HCV clones in culture that the use of *in vitro* synthesized RNA transcripts (cRNA) of defined  
10 structure (produced, *e.g.*, using T7 or SP6 *in vitro* transcription system) is advantageous to transfection of DNA constructs. Indeed, direct transfection of cRNA avoids the involvement of the cell nucleus and therefore potential problems associated with transcriptional regulation, splicing, incorrect 5' and 3'-end processing, and nucleo-cytoplasmic transport (Dash *et al.*, *Am J. Pathol.*, 1997, 151:363-373).

15 However, in contrast to many other (+) strand RNA viruses, construction of self-replicating subgenomic HCV clones in culture turned out to be very difficult. Until recently, only a few successful attempts were reported, each lacking important controls and/or being somewhat controversial. Thus, Baumert *et al.* (*J. Virol.*, 1998, 72:3827-3836) described assembly of poorly characterized HCV-like particles in insect cells upon introduction of a  
20 recombinant baculovirus containing the cDNA encoding HCV ORF. Dash *et al.* (*supra*) and Yoo *et al.* (*J. Virol.*, 1995, 69:32-38) reported successful replication of putative HCV cRNA upon transfection in the human hepatoma cell lines Huh-7 and HepG2, respectively. However, both studies appear to be highly questionable as they describe the propagation of truncated HCV genomes lacking the authentic 3' NTR, which is essential for replication *in*  
25 *vivo* (Yanagi *et al.*, 1999, *supra*; Kolykhalov *et al.*, 2000, *supra*).

The most convincing evidence of a functional *in vitro* cell-based system for replication of recombinant HCV came from a recent report by Lohmann *et al.* (*Science*, 1999, 285:110-113). These authors have described selectable subgenomic HCV RNA molecules replicating after transfection into the human hepatoma cell line Huh-7. Similar subgenomic  
30 HCV replicons capable of propagating in tissue culture are disclosed in PCT Application No. WO 98/39031. Based on the assumption that high expression levels of the structural proteins might be cytotoxic (Moradpour *et al.*, *Biochem. Biophys. Res. Comm.*, 1998, 246:920-924) and the observation that for several (+) strand RNA viruses (*e.g.*, alpha-, flavi- and pestiviruses) the structural proteins are not required for RNA replication (Behrens *et al.*, *J.*

5 Virol., 1998, 72:2364- 2372; Khromykh and Westaway, J. Virol., 1997, 71:1497-1505; Liljestrom and Garoff, Biotechnol., 1991, 9:1356- 1361), Lohmann *et al.* deleted the sequences of the structural proteins in their HCV-derived constructs. In addition, to allow selection for only those cells which support efficient HCV replication, the gene encoding the neomycin phosphotransferase (*neo*) and conferring resistance to the antibiotic G418, was  
10 introduced downstream of the HCV IRES (Figure 3). A second IRES element was included to allow translation of the HCV NS proteins. Upon transfection of these bicistronic RNAs and selection in the presence of G418, only cells supporting replication of HCV-derived RNAs amplified the *neo* gene and developed resistance, whereas non-transfected cells and cells unable to support replication died. The selected cells carried large amounts of HCV  
15 RNAs detectable by Northern blot, or after metabolic radiolabeling with <sup>3</sup>[H]uridine, providing formal proof that these RNAs were actively replicating in the cells. As expected for a replicative intermediate, (-) strand RNA was present in approximately 10-fold lower amounts compared to (+) strand RNA. Finally, HCV proteins could be detected by immunoprecipitation after metabolic radiolabelling with <sup>35</sup>[S]methionine or Western blot and  
20 were confined to the cytoplasm (*see also* Bartenschlager and Lohmann, 2000, *supra*).

Although replicon titers observed in several clones generated by Lohmann *et al.* were several orders of magnitude higher compared to previously available cell culture-based HCV infection systems (described above, *see also* Blight and Gowans, Viral Hepatitis Rev., 1995, 1:143-155), the overall efficiency of clone generation upon transfection was very low. Taken  
25 together with the observation that the majority of cell clones containing replication-competent subgenomic HCV RNAs were growing much slower than the naïve Huh-7 cells or the cells transformed with replication-deficient RNA, it can be concluded that the recombinant HCV replicons of Lohmann *et al.* are toxic to their host cells. Clearly, such cytotoxic replicons are deficient as an *in vitro* model of HCV propagation as they cannot  
30 attain their maximal replication titers and therefore cannot provide sufficient amounts of subgenomic viral nucleic acids required for generation of recombinant anti-viral vaccines, for studies of some of the viral processes, and for sensitive large-scale anti-viral drug screening.

In view of the above, despite the progress made in the last several years, the field is still lacking an efficient stable cell culture system for high titer propagation of recombinant

- 5 HCV. Such a system is needed for (i) further studies of intracellular viral processes (*e.g.*, analysis of HCV receptor binding, cellular infection, replication, virion assembly, and release); (ii) generation and testing of anti-viral vaccines; (iii) screening and testing of anti-viral drugs; (iv) development of targets and methods for HCV diagnostics, and (v) production of concentrated virion and protein stocks (*e.g.*, for structural analysis of virion components  
10 leading to epitope determination for immunotherapy).

The present invention addresses these and other needs in the art by providing novel mutated recombinant HCV-derived nucleic acids and novel rapidly growing "adapted" cell clones supporting their efficient replication.

### SUMMARY OF THE INVENTION

- 15 The present invention provides novel recombinant hepatitis C (HCV)-derived nucleic acids. Preferred subgenomic HCV replicons of the invention include HCVR 2 (SEQ ID NO: 2), HVCR 8 (SEQ ID NO: 3), HCVR 9 (SEQ ID NO: 4), HCVR 22 (SEQ ID NO: 5) and HCVR 24 (SEQ ID NO: 6). These replicons are derived from the parental HCV genotype 1b-based recombinant clone I377-NS3-3'UTR (SEQ ID NO: 1) and contain multiple  
20 nucleotide changes (*e.g.*, as shown in Table I, *see* Example 1) which occurred following their prolonged replication in the Huh-7-derived cell clones of the present invention under stringent selection conditions. At least some of these mutations are indicative of the non-structural HCV genome regions which are responsible for high-titer viral replication and virus-induced cytotoxicity.

- 25 In a specific embodiment, the present invention is directed to plasmid clones which can be transcribed to produce self-replicating recombinant HCV RNAs of the invention.

- In a separate embodiment, the instant invention includes a method for generating novel efficiently replicating recombinant HCV-derived nucleic acids containing the critical elements of the HCV replicons described above and comprising from 5' to 3' on the  
30 positive-sense nucleic acid (1) a functional 5' non-translated region (NTR) comprising an extreme 5'-terminal conserved sequence, (2) at least one open reading frame (ORF) encoding a heterologous gene operatively associated with an expression control sequence, wherein the

5 heterologous gene and expression control sequence are oriented on the positive-strand nucleic acid molecule, (3) an ORF encoding at least a portion of an HCV polyprotein whose cleavage products form functional components of HCV virus particles and RNA replication machinery, and (4) a 3' NTR comprising an extreme 3'-terminal conserved sequence. In a specific embodiment, where the selection marker ORF is a drug resistance gene, this gene is a  
10 neomycin resistance gene (*neo*) operatively associated with an internal ribosome entry site (IRES).

The instant invention also includes a method for propagating the disclosed recombinant HCV-derived nucleic acids *in vitro* by culturing a cell line transfected or infected with an appropriate amount of HCV RNA, *e.g.*, as produced from the plasmid clones  
15 recited above, under conditions that permit replication of the HCV RNA. In a specific embodiment, replication of the disclosed novel HCV-derived nucleic acids in susceptible cultured cells leads to the generation of potentially infectious recombinant viral particles which can be used as an attenuated anti-HCV vaccine.

Accordingly, in conjunction with the recombinant nucleic acids disclosed herein, the  
20 present invention advantageously provides cell lines, which are susceptible to HCV infection and/or transfection and support replication of such recombinant nucleic acids. In a preferred embodiment, the susceptible cell line of the invention is a human hepatoma cell line Huh-7.

The invention further provides stable "adapted" cell clones which are derived from  
25 Huh-7 cell line and are characterized by the growth properties which are similar to or indistinguishable from the naive (*i.e.*, untransfected) Huh-7 cells (*e.g.*, as shown in Figure 5). According to the instant invention, these "adapted" cell clones are able to support efficient replication of subgenomic HCV RNAs. Some of the most efficiently propagating cell clones of the present invention, *i.e.*, HCVR 2, 8, 9, 22 and 24, were deposited with the American  
30 Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209, USA on September 20, 2000, and assigned Accession Nos. PTA-2489, PTA-2490, PTA-2486, PTA-2487, and PTA-2488, respectively.

The materials disclosed herein provide methods for screening (both *in vitro* and *in vivo*) for agents capable of modulating HCV infection and/or replication and/or virion

5 assembly. Such methods include administering a candidate agent to HCV-replicating cell line(s) of the invention, and testing for an increase or decrease in a level of subgenomic HCV replication or HCV-associated protein expression compared to a level of HCV replication or HCV-associated protein expression in a control cell line transfected with replication-defective construct or in the same cell line prior to administration of the candidate agent, wherein a  
10 decrease in the level of HCV replication or HCV-associated protein expression is indicative of the inhibitory activity of the agent.

In a specific embodiment, HCV-replicating cell lines of the present invention provide a convenient system for high-throughput initial screening of potential anti-HCV therapeutics.

Further provided herein is a method for generating massive quantities of recombinant  
15 HCV replicons (which can serve as a basis for anti-HCV vaccine development) from the cell clones of the present invention, said replicons being produced either as intracellular nucleic acids or as infectious or non-infectious recombinant viral particles.

The present invention also has significant diagnostic implications. For example, the invention provides an *in vitro* method for detecting antibodies to HCV in a biological sample  
20 from a subject comprising contacting the sample with HCV-replicating cells, cellular fractions, isolated HCV-derived proteins, or HCV-derived viral particles prepared as described above. The contacting operation is conducted under conditions that permit binding of HCV-specific antibodies in the sample to the HCV protein(s); and detecting binding of antibodies in the sample to the HCV-derived protein(s). Detecting binding of antibodies in  
25 the sample to the HCV protein(s) is indicative of the presence of HCV infection in the subject from which the sample was derived.

In summary, the present invention provides nucleic acids encoding recombinant HCV replicons, which are capable of efficient propagation and expression of HCV-derived proteins in a cell culture system.

30 The invention further provides susceptible cell lines (and "adapted" rapidly growing cell clones derived from them), which support high titer replication of recombinant HCV-derived nucleic acids.

By providing cell clones supporting efficient subgenomic HCV replication, the present invention provides (i) *in vitro* cell culture models of HCV propagation; (ii) systems



5 for screening candidate anti-viral compounds and evaluating drug resistance; (iii) methods for diagnosing HCV infection, and (iv) systems for production of large quantities of HCV-derived components or recombinant viral particles for antibody generation and/or vaccine development.

The present invention meets these and other objects of the invention, as set forth in  
10 greater detail in the Detailed Description and Examples, including the accompanying Drawings.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

**FIGURE 1 (prior art)** depicts HCV genome structure and polyprotein processing. At the top is a schematic representation of the viral genome with the structural and nonstructural protein coding regions, and the 5' and 3' NTRs. Boxes below the genome  
15 identify proteins generated by the proteolytic processing cascade. Different types of arrows represent different types of proteases responsible for polyprotein processing at each particular site.

**FIGURE 2 (prior art)** is a schematic representation of the HCV life cycle  
20 comprising the following stages: (1) penetration of the host cell and liberation of the (+) strand genomic RNA (cRNA) from the virus particle into the cytoplasm; (2) translation of the input cRNA, processing of the polyprotein and formation of a replicase complex associated with intracellular membranes; (3) utilization of the input (+) strand for synthesis of a (-) strand RNA intermediate; (4) production of new (+) strand RNA molecules which can be  
25 used for synthesis of new (-) strands, for polyprotein expression or for packaging into progeny virions; (5) release of virus from the infected cell via cellular secretion pathway resulting in formation of cell-derived viral membrane envelope.

5           **FIGURE 3** is a schematic representation of the recombinant parental HCV replicon I377/NS3-3'UTR (SEQ ID NO: 1) composed of the 5'HCV-IRES, the *neo* gene, the EMCV-IRES, and HCV sequences from NS3 up to the authentic 3' end.

**FIGURE 4** depicts detection of (+) strand HCV RNA in subpassaged Huh-7 cell clones by RT-PCR using primers flanking the region corresponding to NS5B nt 7435-7750.  
10   Shown is 2% agarose gel electrophoresis analysis of RT-PCR fragments obtained by amplifying total RNA isolated from G418-resistant Huh-7 cell clones HCVR 2, 6, 8, 9 and 12 (lanes 2-6). As a negative control, PCR was performed without a DNA template (lane 1). Lane MW shows position of molecular size markers.

**FIGURE 5** is a graphic representation of relative growth rates of Huh-7-derived  
15   "adapted" cell clones supporting efficient subgenomic HCV replication. Growth rates (shown as a function of optical density after sulphorhodamine staining for total cellular proteins) of clones HCVR 2, 8, 9, 22 and 24 are compared to the growth rate of naïve non-transfected Huh-7 cells (black line) at 9 weeks post-transfection.

#### **DETAILED DESCRIPTION OF THE INVENTION**

20           All patent applications, patents, and literature references cited herein are hereby incorporated by reference in their entirety.

          The present invention provides novel recombinant hepatitis C (HCV) nucleic acids containing all HCV non-translated sequences required for replication and expression, all or a portion of the HCV ORF, as well as one or more heterologous genes. In a preferred  
25   embodiment, disclosed herein are recombinant HCV-derived nucleic acids (e.g., HCVR 2 [SEQ ID NO: 2], HCVR 8 [SEQ ID NO: 3], HCVR 9 [SEQ ID NO: 4], HCVR 22 [SEQ ID NO: 5] and HCVR 24 [SEQ ID NO: 6]) which differ from the parental chimeric HCV replicon used for their generation (I377/NS3-3'UTR, SEQ ID NO: 1) in a number of positions

5 located at various parts of the replicon genome (*see, e.g.*, Table I). At least some of these mutant subgenomic HCV nucleic acids have a much lower cytotoxicity, as evident from their growth potential (Figure 5) and, subsequently, have a significantly higher rate of productive transfection compared to the previously described recombinant HCV replicons. Due to these and other advantageous properties, the subgenomic HCV nucleic acids of the present  
10 invention are capable of efficient high titer replication in the susceptible cell lines of the present invention.

In conjunction with HCV-derived nucleic acids, the present invention further provides susceptible cell lines and, in particular, novel "adapted" rapidly growing cell clones derived from human hepatoma cell line Huh-7 (*e.g.*, clones HCVR 2, 8, 9, 22 and 24), said cell lines  
15 and "adapted" clones being capable of supporting efficient replication of subgenomic recombinant HCV RNAs.

Another embodiment is a sensitive high-throughput method for screening anti-HCV therapeutics by putting them in contact with the cell clones of the present invention and determining their effect on propagation of subgenomic HCV replicons.

20 Further provided herein is a method for generating large quantities of recombinant HCV replicons from the cell clones of the present invention. These replicons can be generated for the purpose of providing a recombinant attenuated anti-HCV vaccine and can be produced either as intracellular nucleic acids or as infectious or non-infectious recombinant viral particles.

25 By providing cell clones supporting efficient subgenomic HCV replication, the present invention provides (i) *in vitro* cell culture models of HCV propagation; (ii) systems for screening candidate anti-viral compounds and evaluating drug resistance; (iii) methods for diagnosing HCV infection, and (iv) systems for production of large quantities of HCV-derived components or recombinant viral particles for antibody generation and/or vaccine  
30 development.

### Definitions

5       A "nucleic acid molecule" refers to the phosphate ester polymeric form of ribonucleosides (adenosine, guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine, deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules"), in either single stranded form, or a double-stranded form. Double stranded DNA-DNA, DNA-RNA and RNA-RNA duplexes are possible. The term nucleic  
10   acid molecule, and in particular DNA or RNA molecule, refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. In discussing the structure of particular nucleic acid molecules, sequences or regions may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction. A "recombinant DNA molecule" is a DNA molecule that has undergone a  
15   molecular biological manipulation.

      A "coding sequence" or a sequence "encoding" an expression product, such as a RNA, polypeptide, protein, or enzyme, is a nucleotide sequence that, when expressed, results in the production of that RNA, polypeptide, protein, or enzyme, *i.e.*, the nucleotide sequence encodes an amino acid sequence for that polypeptide, protein or enzyme. In the present  
20   invention, translation of HCV-derived (+) strand RNA initially yields a polyprotein, which is cleaved during post-translational processing to yield functional viral proteins.

      The term "gene" means a DNA sequence that codes for or corresponds to a particular sequence of amino acids, which comprise all or part of one or more proteins or enzymes.

      A coding sequence is "under the control" or "operatively associated with"  
25   transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then translated into the protein encoded by the coding sequence.

      The terms "express" and "expression" mean allowing or causing the information in a gene or nucleic acid sequence to become manifest, for example producing a protein by  
30   activating the cellular functions involved in transcription and translation of a corresponding gene or nucleic acid sequence. A nucleic acid sequence is expressed in or by a cell to form

5 an "expression product" such as a mRNA or a protein. The expression product itself, *e.g.* the resulting protein, may also be said to be "expressed" by the cell.

The term "polypeptide" refers to a polymer of amino acids and does not refer to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not refer to, or exclude, post-translational  
10 modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations, and the like.

As used herein, the term "conservative mutation" or "conservative nucleotide change" is used to define a nucleotide change, which occurs with high frequency within quasispecies. "Sequence-conservative variants" are those in which a change of one or more nucleotides in a  
15 given codon position results in no alteration in the amino acid encoded at that position. "Function-conservative variants" are those in which a given amino acid residue in a protein or enzyme has been changed without altering the overall conformation and function of the polypeptide, including, but not limited to, replacement of an amino acid with one having similar properties (such as, for example, polarity, hydrophobicity, size of the side chain,  
20 hydrogen bonding potential, and the like).

The term "hepatitis C virus" or "HCV" is used herein to define a viral species of which pathogenic strains cause hepatitis C, also known as non-A, non-B hepatitis.

As used herein, the term "viral RNA", which includes HCV RNA, refers to RNA from the viral genome, fragments thereof, transcripts thereof, and mutant sequences derived  
25 therefrom.

As used herein, a "(+) stranded genome" or a "positive-stranded genome" of a virus is one in which the genome, whether RNA or DNA, is single-stranded and encodes a viral or virus-derived polypeptide(s). Examples of positive-stranded RNA viruses include Togaviridae, Coronaviridae, Retroviridae, Picornaviridae, and Caliciviridae. Included also,  
30 are the Flaviviridae, which were formerly classified as Togaviridae (*see* Fields and Knipe, *Fundamental Virology*, Raven Press, 1986).

As used herein, a "replicative intermediate" of an HCV genome or a "(-) stranded genome" or a "negative-stranded genome" refers to an RNA strand or fragment thereof, which is complementary to the viral genome, and which is synthesized during the course of

5 viral replication; the replicative intermediate functions as a template for the synthesis of (+) RNA strands.

As used herein, "purified HCV virions" refers to a preparation of HCV viral or virus-like particles that have been isolated from the cellular constituents with which the virus normally associates, and from other types of viruses that may be present in the infected  
10 tissue. The techniques for isolating viruses are known to those of skill in the art, and include, for example, centrifugation and affinity chromatography.

A "construct" is a chimeric virus or nucleic acid encoding a chimeric virus, such as positive viral genomic RNA or a DNA that can be transcribed to produce viral genomic RNA.

15 The term "chimeric" is used herein in its usual sense: a construct or protein or virus resulting from the combination of genes from two or more different sources, in which the different parts of the chimera function together. The genes are fused, where necessary in-frame, in a single genetic construct. As used herein, the term "chimeric" refers specifically to recombinant HCV-derived nucleic acids or proteins or virions.

20 The term "chimeric virus genome" or "recombinant virus" or "subgenomic HCV replicon" as used herein refers to the genome of the HCV that is modified by insertion or substitution of sequences. In some instances, the virus-derived replicon which later undergoes additional changes (*e.g.*, as a result of *in vitro* manipulations or *in vivo* selection) may be referred to as a "parent" genome or replicon. In general, according to the present  
25 invention, the recombinant virus genome will include various parts of the parent virus genome, said parts comprising without limitation genes encoding proteins involved in replication, infectivity, tropism, and life cycle.

The term "junction site" is used herein to refer to the amino acid sequence joining two different proteins of a virus-derived polyprotein that is recognized and cleaved by a protease,  
30 *e.g.*, HCV NS3. Various HCV NS3 junction sites are known, including the NS3/NS4A, NS4A/NS4B, NS4B/NS5A, and NS5A/NS5B junction sites. Any of these can be used to substitute for an endogenous junction site of the virus used in the chimeric construct.

5           The term "internal ribosome entry site" or "IRES" defines a special region of some viral and cellular mRNA molecules that is capable of cap-independent binding of the ribosome to initiate translation.

          As used herein, "infectious" refers to the ability of a virus to enter and replicate in a cell and to produce viral particles. Infectivity can be evaluated either by detecting virus, *i.e.*,  
10   viral load, or by observing disease progression in an animal. Virus (viral load) can be detected by the presence of viral (+) strand RNAs and/or (-) strand replication intermediates, *e.g.*, detected by RT-PCR or direct hybridization techniques. It can also be detected, if present in sufficient amount, by the presence of replicon-derived proteins, *e.g.*, detected by immunoassay or biochemical techniques. In another alternative, a culture medium isolated  
15   from a cell line supporting viral replication or extracts/samples from an animal are used to infect naive cells in culture. The important aspects of the determination of viral infectivity *in vivo* (*i.e.*, in infected subjects) is the development of either an acute or chronic viral infection, which, in turn, may include either overt pathology or merely replication and propagation of the virus.

20           The term "viral load" or "viral titer" is used herein to refer to a quantitative amount of virus in a cell culture or in an infected animal. The term "titer" can be also used to refer to a quantitative amount of virus-derived replicons produced within a susceptible cell. "Disease progression" refers to the course of disease or symptoms of an infected animal, and may include acute or chronic disease symptoms. "Pathogenesis" is a particular indication of a  
25   disease progression, and refers to the pathogenic effects of viral infection, including morbidity and mortality. In the present invention, main pathologic effects of HCV are observed in the hepatic tissue.

          An "individual" or "subject" or "animal", as used herein, refers to vertebrates, particularly members of the mammalian species and includes, but is not limited to, rodents,  
30   rabbits, shrews, and primates, the latter including humans.

          As used herein, a "biological sample" refers to a sample of tissue or fluid isolated from an individual, including but not limited to, for example, plasma, serum, spinal fluid, lymph fluid, the external sections of the skin, respiratory, intestinal and genitourinary tracts, tears, saliva, milk, blood cells, tumors, organs, and also samples of *in vitro* cell culture

5 constituents (including, but not limited to, conditioned medium resulting from the growth of cultured cells, putatively viral infected cells, recombinant cells, and cell components).

As used herein, the term "quasispecies" means a collection of microvariants of a predominant HCV genome sequence ("master sequence"), said microvariants being formed in a single infected animal or even in a single cell clone as a result of high mutation rate  
10 during HCV replication.

As used herein, an "*in vitro* cell system" or an "extracorporeal cell system" refers to cells which are replicated outside of the body, *i.e.*, cell systems not found in nature; as such, the term includes primary cultures and cell lines.

"Primary cultures", as used herein, refers to a culture of cells that is directly derived  
15 from cells or tissues from an individual, as well as cells derived by passage from these cells, but not to immortalized cells.

As used herein, "cell line" refers to a population of cells capable of continuous or prolonged growth and division *in vitro*. The term "cell lines" also includes immortalized cells. Often, cell lines are clonal populations derived from a single progenitor cell. Such cell  
20 lines are also termed "cell clones". It is further known in the art that spontaneous or induced changes can occur in karyotype during storage or transfer of such clonal populations. Therefore, cells derived from the cell clones referred to may not be precisely identical to the ancestral cells or cultures. *According to the present invention, if such cell clones are capable of supporting efficient replication of HCV-derived RNAs without a significant decrease in*  
25 *their growth properties, they are termed "adapted" cell clones.*

The term "host cell" means any cultured cell or any cell of any organism that is susceptible to infection by or propagation of a wild-type HCV or a chimeric virus construct of the invention. Host cells can further be used for screening or other assays, as described herein. A potentially susceptible cell that has not been transfected or infected with virus-  
30 derived nucleic acids is termed a "naive" cell.

The term "transfection" means the introduction of a foreign nucleic acid into a cell. The introduced gene or sequence may also be called a "cloned" or "foreign" gene or sequence, may include regulatory or control sequences, such as start, stop, promoter, signal, secretion, or other sequences used by a cellular molecular machinery. A host cell that



5 receives and expresses introduced DNA or RNA has been "transfected" and is a "transfectant" or a "recombinant cell". According to the present invention, if a transfection of HCV-derived nucleic acid results in its subgenomic replication and HCV-derived protein expression, the transfection is termed "productive".

10 The term "antibody", as used herein, includes both monoclonal and polyclonal antibodies. Additionally, single polypeptide chain antigen-binding proteins, see U.S. Pat. No. 4,946,778, are included within the term "antibody".

As used herein, "epitope" refers to an antigenic determinant of a polypeptide; an epitope can comprise 3 or more amino acids in a spatial conformation unique to the epitope. Generally an epitope consists of at least 5 such amino acids and, more usually, consists of at  
15 least 8-10 such amino acids. Methods of determining the spatial conformation of amino acids are known in the art, and include, for example, x-ray crystallography and nuclear magnetic resonance (NMR).

"Immunologically identifiable with/as" refers to the presence of epitope(s) and polypeptides(s) that are also present in the designated polypeptide(s), herein usually HCV  
20 proteins. Immunological identity may be determined by antibody binding and/or competition in binding; these techniques are known to those of average skill in the art.

A polypeptide is "immunologically reactive" with an antibody when it binds to an antibody due to antibody recognition of a specific epitope contained within the polypeptide. Immunological reactivity may be determined by antibody binding, more particularly by the  
25 kinetics of antibody binding, and/or by competition in binding using as competitor(s) a known polypeptide(s) containing an epitope against which the antibody is directed. The techniques for determining whether a polypeptide is immunologically reactive with an antibody are known in the art.

As used herein, the term "immunogenic polypeptide" refers to a polypeptide that  
30 elicits a cellular and or humoral immune response, whether alone or when linked to a carrier, in the presence or absence of an adjuvant.

In a specific embodiment, the term "about" or "approximately" means within a statistically meaningful range of a value. Such a range can be within an order of magnitude, preferably within 50%, more preferably within 20%, more preferably still within 10%, and

5 even more preferably within 5% of a given value or range. The allowable variation encompassed by the term "about" or "approximately" depends on the particular system under study, and can be readily appreciated by one of ordinary skill in the art.

### General Techniques

10 In accordance with the present invention there may be employed conventional molecular biology, microbiology, biochemistry, genetics and immunology techniques within the skill of the art for the production of recombinant HCV nucleic acids, HCV-replicating cell cultures, infectious viral particles, viral and recombinant proteins, antibodies, etc. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch and Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition, 1989, Cold Spring Harbor Laboratory Press (herein "Sambrook et al., 1989"); *DNA Cloning: A Practical Approach*, 15 Glover ed., Volumes I and II, 1985; *Oligonucleotide Synthesis*, Gait ed., 1984; *Nucleic Acid Hybridization*, Hames and Higgins eds., 1985; *Transcription And Translation*, Hames and Higgins eds., 1984; *Animal Cell Culture*, Freshney ed., 1986; *Immobilized Cells And Enzymes*, IRL Press, 1986; Perbal, *A Practical Guide To Molecular Cloning*, 1984; *Current Protocols in Molecular Biology*, Ausubel et al., eds., 1994, John Wiley & Sons.

### Construction of Recombinant HCV Replicons

20 In a broad aspect, the present invention is directed to genetically engineered HCV nucleic acid clones which comprise from 5' to 3' on the positive-sense nucleic acid (1) a functional 5' non-translated region (NTR) comprising an extreme 5'-terminal conserved sequence, (2) at least one open reading frame (ORF) encoding a heterologous gene operatively associated with an expression control sequence, wherein the heterologous gene and expression control sequence are oriented on the positive-strand nucleic acid molecule, (3) 25 an ORF encoding at least a portion of an HCV polyprotein whose cleavage products form functional components of HCV virus particles and RNA replication machinery, and (4) a 3' NTR comprising an extreme 3'-terminal conserved sequence. According to the present invention, the heterologous gene can be a drug resistance gene or a reporter gene. In a

5 specific embodiment, where the selection marker ORF is a drug resistance gene, this gene is a neomycin resistance gene (*neo*) operatively associated with an internal ribosome entry site (IRES). Alternatively, the heterologous gene can be a therapeutic gene, or a gene encoding a vaccine antigen, *i.e.*, for gene therapy or gene vaccine applications, respectively.

10 In a preferred embodiment, the subgenomic HCV replicons are derived from a "parent" HCV genotype 1b-based chimeric replicon I377/NS3-3'UTR (SEQ ID NO: 1, see also Figure 3). Due to their prolonged propagation under stringent selection conditions (*e.g.*, 1mg/ml G418, *see Example 1, infra*) in susceptible cell lines, the subgenomic HCV replicons of the present invention have acquired multiple mutations. Accordingly, also disclosed herein are positions of at least some of the nucleotide changes identified in these mutant  
15 HCV replicons (*e.g.*, as shown in Table I), said nucleotide changes being indicative of the genome regions which are responsible for high-titer virus replication and for virus-induced cytotoxicity.

The actual molecular biological techniques required to generate fusion between heterologous sequences and specific fragments of HCV genome are routine. For example, a  
20 recombinant replicon can be constructed using PCR (*see Lohmann et al., supra; Example 1, infra*).

Naturally, as noted above, the HCV nucleic acids of the invention are selected from the group consisting of double stranded DNA, (+) strand cDNA, or (-) strand DNA, or (+) strand RNA or (-) strand RNA. Thus, where particular sequences of nucleic acids of the  
25 invention are set forth, both DNA and corresponding RNA are intended, including positive and negative strands thereof.

According to the instant invention, an HCV DNA may be inserted in a plasmid vector for transcription of the corresponding HCV RNA. Thus, the HCV DNA may comprise a promoter 5' of the 5'-NTR on (+) strand DNA, whereby transcription of template DNA from  
30 the promoter produces replication-competent RNA. The promoter can be a member selected from the group consisting of a eukaryotic promoter, yeast promoter, plant promoter, bacterial promoter, and viral promoter. The preferred promoter is phage T7 promoter (*see Example 1, infra*).

5 In a specific embodiment, the present invention is directed to a plasmid clone, pAn/HCVR1, which can be transcribed to produce self-replicating recombinant "parental" HCV RNA transcript I377/NS3-3'UTR (*see* Figure 3).

Also disclosed herein are cRNA molecules transcribed from the recombinant HCV-derived DNA plasmid clones set forth above. These cRNA molecules are used for  
10 transfection into the cell lines of the present invention.

The present invention further advantageously provides DNA plasmid vectors containing inserts derived from the mutant RNA replicons disclosed herein using reverse transcription followed by PCR-amplification or other molecular biology techniques well known in the art.

15 Naturally, the invention also includes derivatives of all of the disclosed nucleic acids, selected from the group consisting of derivatives produced by substitution of homologous regions from other HCV isolates or genotypes; derivatives produced by mutagenesis; derivatives selected from the group consisting of infectious, adapted, live-attenuated, replication competent non-infectious, and defective variants; derivatives comprising  
20 additional heterologous gene(s) operatively associated with an expression control sequence; and derivatives consisting of a functional fragment of any of the above-mentioned derivatives. Alternatively, portions of the disclosed nucleic acids, such as the 5' NTRs, the polyprotein coding regions, the 3'-NTRs or more generally any coding or non-translated region of the HCV genome, can be substituted with a corresponding region from a different  
25 HCV genotype to generate a new chimeric clone, or by extension, clones of other isolates and genotypes. For example, an HCV-1a or -2a polyprotein coding region (or consensus polyprotein coding regions) can be substituted for the HCV-1b polyprotein coding region of the deposited clones.

By providing the engineered HCV nucleic acids of the invention, the present  
30 inventors have made it possible to dissect the HCV replication machinery and protein activity. The instant invention has also made possible the preparation of various HCV derivatives such as attenuated, expressing heterologous gene(s), replication-competent non-infectious, replication-defective infection-competent, and replication-defective non-

- 5 infectious. The recombinant HCV DNA clones or RNAs of the invention can also be used in numerous methods, or to derive authentic HCV components, as set forth below.

### **Production of "Adapted" Cell Clones Supporting Efficient Subgenomic HCV Replication**

The instant invention further provides a method for propagating the disclosed recombinant HCV-derived nucleic acids *in vitro* comprising culturing a cell line transfected or infected with an appropriate amount of HCV RNA, *e.g.*, as produced from the plasmid clones recited above, under conditions that permit replication of said HCV-derived RNA.

By providing novel recombinant HCV nucleic acids, in particular, nucleic acids containing mutations in the regions responsible for cytotoxicity and/or replicon titer, the present invention provides a method for generating stable cell clones supporting efficient replication of subgenomic HCV RNAs.

15 Accordingly, in conjunction with the recombinant nucleic acids disclosed herein, the present invention advantageously provides susceptible stable cell lines which, upon transfection or infection, can support high titer replication of the disclosed HCV-derived nucleic acids. The susceptible cell lines of the invention include without limitation human hepatoma cell lines Huh-7, HepG2, and PH5CH; *T. belangeri* liver cell line MBTL; human  
20 diploid fibroblast cell line VERO; secondary monkey kidney cell line CV-1; T cell lines MT-2, HPBMA10-2, and MOLT-4, and B cell line Daudi.

As disclosed herein, the HCV-replicating stable cell lines of the present invention are characterized by the growth rates which are not less than 10% of the growth rate of the corresponding naïve (non-transfected) cell lines. In a preferred embodiment, these growth  
25 rates are not less than 25% of the growth rate of the corresponding naïve (non-transfected) cell lines, or, even more preferably, not less than 90% of the growth rate of the corresponding naïve (non-transfected) cell lines.

In a preferred embodiment, the susceptible cell line is the human hepatoma cell line Huh-7. As disclosed herein, the "adapted" cell clones derived from Huh-7 cell line support  
30 efficient replication of subgenomic HCV nucleic acids of the present invention and are

5 characterized by the growth properties which are similar to or indistinguishable from the  
naive (*i.e.*, non-transfected) Huh-7 cells (*see, e.g.*, Figure 5). Besides promoting the  
generation of novel quasispecies of recombinant replicons (*e.g.*, containing mutations leading  
to lowered cytotoxicity and increased replication efficiency) these "adapted" cell clones  
10 might have acquired certain mutations in their own genomes, said mutations distinguishing  
them from the parental naive cells and leading to their improved ability to support replication  
of subgenomic HCV RNAs. Some of the most efficiently propagating cell clones of the  
present invention, *i.e.*, HCVR 2, 8, 9, 22 and 24, have been deposited with the American  
Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209,  
USA on September 20, 2000, and assigned Accession Nos. PTA-2489, PTA-2490, PTA-  
15 2486, PTA-2487, and PTA-2488, respectively.

The methods for detection of subgenomic HCV replicons in the cell lines of the  
present invention are well known in the art and include without limitation (i) Northern blot or  
strand-specific reverse transcriptase-polymerase chain reaction (RT-PCR) detection of (-)  
strand replicative RNA intermediates; (ii) determination of an increase of (+) strand RNA  
20 using either b-DNA assay or quantitative RT-PCR, (iii) detection of expression of a  
heterologous reporter gene encoded by a recombinant HCV replicon (*e.g.*, by assaying  
fluorescence, luminescence or enzymatic activity) and (iv) detection of HCV-derived proteins  
(*e.g.*, by Western blot or immuno-fluorescence).

The permissive cell lines that are identified using the nucleic acids of the invention  
25 are very useful, *inter alia*, for studying the HCV infection and propagation, isolating  
functional components of HCV, and for sensitive, fast diagnostic and therapeutic  
applications.

In a specific embodiment, the permissive cell line and the "adapted" cell clones of the  
present invention are used to select novel mutant recombinant HCV replicon quasispecies.  
30 By providing HCV nucleic acids comprising a selectable marker, the present invention  
provides a method for selecting novel mutated chimeric HCV replicons characterized by low  
cytotoxicity, high productive transformation efficiency, and high replicon titer attained; said  
method comprising (i) transformation of a susceptible cell line followed by (ii) prolonged  
culturing under stringent selection conditions which allow only the survival of cells

- 5 supporting high titer HCV replication, and (iii) isolation of recombinant replicons from the cell clones characterized by the best growth properties. Methods for identification of the novel mutations contained within these selected replicons are well known in the art and include, for example, a sequence of steps: reverse transcription -> PCR amplification -> automated sequencing -> computer sequence analysis.

### Production of Recombinant Virions

- 10 The invention further provides a method for producing recombinant HCV virus-like particles, comprising isolating HCV virus-like particles from the cell lines of the invention (or their culture medium) under conditions that permit subgenomic HCV replication and virus particle formation. The present invention extends to a recombinant HCV-derived virus particle comprising a replication-competent subgenomic HCV RNA, or a  
15 replication-defective HCV-derived RNA; said recombinant HCV virus particle being either infection-competent or infection-defective. In a specific embodiment the recombinant viral particles produced in the cell lines of the present invention provide an attenuated recombinant vaccine to be administered to an individual to produce an anti-viral immune response. Alternatively, isolated HCV-derived proteins expressed in the cell lines of the present  
20 invention can represent starting materials for an HCV vaccine. Preferably, a vaccine of the invention includes a pharmaceutically acceptable adjuvant. Representative but not limiting examples of adjuvants include Complete and Incomplete Freund's Adjuvant, detoxified endotoxins, mineral oils, surface active substances such as lipolecithin, pluronic polyols, polyanions, peptides, oil or hydrocarbon emulsions, immunostimulatory compounds, *Bacille*  
25 *Calmette-Guerin* (BCG), and the synthetic adjuvant QS-21 (McCune *et al.*, Cancer, 1979; 43:1619).

### Diagnostic and Therapeutic Applications

By providing for insertion of heterologous genes in the recombinant HCV nucleic acids, the present invention provides a method for transducing an animal susceptible to HCV infection with said heterologous genes by administering an amount of the HCV RNA to the

5 animal either directly or in the form of infectious recombinant viral particles produced in the cell lines disclosed herein. HCV-mediated introduction of heterologous genes in susceptible animals can be useful, for example, for gene therapy or gene vaccination which is targeted to hepatic tissues.

Also provided is an *in vitro* cell-free assay system for HCV replication comprising  
10 HCV-derived template cRNA of the invention, *e.g.*, as transcribed from a plasmid of the invention as set forth above, functional HCV replicase components, and an isotonic buffered medium comprising ribonucleotide triphosphate bases. These elements provide the replication machinery and raw materials (NTPs).

In a further embodiment, the invention provides a method for producing novel  
15 polyclonal antibodies to HCV-derived proteins and/or recombinant viral particles comprising administering an immunogenic amount of HCV-derived proteins isolated from the cell cultures or the *in vitro* cell-free system described above to an animal, and isolating generated anti-HCV antibodies. A further method for producing antibodies to HCV comprises screening a human antibody library for reactivity with HCV-derived proteins of the invention  
20 and selecting a clone from the library that expresses a reactive antibody. Alternatively, novel monoclonal anti-HCV antibodies can be produced in hybridoma cell lines using techniques well known in the art.

According to the present invention, the usefulness of newly generated antibodies can be assayed by measuring their affinity and specificity (*e.g.*, upon their application to isolated  
25 HCV-derived antigens and/or HCV replicating cell lines of the invention and/or liver tissue sections from chronically infected animals).

The novel anti-HCV antibodies disclosed herein may be used diagnostically, *e.g.*, to detect the presence and/or propagation of HCV in a cell culture or in an animal. Alternatively, these antibodies may be used therapeutically, *e.g.*, in passive immunotherapy.

30 The present invention further advantageously provides methods for screening (both *in vitro* and *in vivo*) for agents capable of modulating HCV infection and/or replication and/or virion assembly. Such methods include administering a candidate agent to HCV infected or transfected cell line(s) of the invention, and testing for an increase or decrease in a level of subgenomic HCV replication or HCV-associated protein expression compared to a level of



5 HCV replication or HCV-associated protein expression in a control cell line transfected with replication-defective construct or in the same cell line prior to administration of the candidate agent, wherein a decrease in the level of HCV replication (*i.e.*, decrease in intra- or extracellular levels of recombinant HCV (+) and/or (-) RNA) or HCV-associated protein expression (*e.g.*, decrease in intra- or extracellular levels of a reporter protein) is indicative of  
10 the inhibitory activity of the agent. Agent-mediated inhibition of virion formation can be detected microscopically (performed directly or after immunostaining); and changes in infectivity of generated HCV virus particles can be assayed by isolating them from the cell culture medium and applying to naive cells or a susceptible animal model.

In a specific embodiment, HCV-replicating cell lines of the present invention provide  
15 a convenient system for high-throughput initial screening of potential anti-HCV therapeutics. Such high-throughput screening system involves applying test compounds to the microcultures of cell clones supporting subgenomic HCV replication (growing, *e.g.*, in 96- or 324-well microtiter plates) followed by measuring changes (*e.g.*, using multi-plate readers or scanners) in HCV replication and/or HCV-associated protein expression and/or HCV  
20 infectivity. According to the instant invention, candidate therapeutic compounds include without limitation small molecule enzyme inhibitors (*e.g.*, helating agents), inhibitory peptides, inhibitory (*e.g.*, transdominant-negative) proteins, antibodies, ribozymes, and antisense nucleic acids.

A further method for screening for agents capable of modulating HCV propagation  
25 involves a cell-free system described above. This method (which can be also performed in a high-throughput format) comprises contacting the *in vitro* cell-free system of the invention with a candidate agent and testing for an increase or decrease in a level of HCV replication or HCV-associated protein expression compared to a level of HCV replication or HCV-associated protein expression in a control cell-based system or a control cell-free system prior  
30 to administration of the candidate agent; wherein a decrease in the level of HCV replication or HCV-associated protein expression compared to the control level is indicative of the ability of the agent to inhibit HCV propagation.

As disclosed herein, the anti-HCV therapeutic compounds identified using the initial *in vitro* screening methods of the present invention can be further characterized for their

5 ability to affect subgenomic HCV propagation using secondary screens in cell cultures and/or susceptible animal models. Based on the tropism of the HCV, a preferred small animal model of the present invention is a tree shrew *Tupaia belangeri chinensis*. A preferred large animal model is a chimpanzee. Test animals will be treated with the candidate compounds that produced the strongest inhibitory effects in cell culture-based assays (control animals  
10 would not be treated, and, if available, a positive control could also be employed). A compound that protects animals from infection by the chimeric virus and/or inhibits viral propagation leading to pathogenicity, would be an attractive candidate for development of an agent for treatment/prevention of HCV infection. In addition, the animal models provide a platform for pharmacokinetic and toxicology studies.

15 The present invention also has significant diagnostic implications. For example, HCV-specific antibodies prepared according to the invention can be used to detect HCV presence and/or propagation in various biological samples. On the other hand, the invention provides an *in vitro* method for detecting antibodies to HCV in a biological sample from a subject comprising contacting said sample with HCV-replicating cells, cellular fractions,  
20 isolated HCV-derived proteins, or HCV-derived viral particles prepared as described above, under conditions that permit interaction of HCV-specific antibodies in the sample with the HCV protein(s), followed by detecting binding of the antibodies in the sample to the HCV-derived protein(s), wherein said binding is indicative of the presence of HCV infection in the subject from which the sample was derived. In the foregoing methods, the biological sample  
25 can be derived without limitation from blood, serum, plasma, blood cells, lymphocytes, or liver tissue biopsy. Techniques for isolating proteins and cellular fractions useful in the foregoing diagnostic methods are also well known in the art.

In a related aspect, the invention also provides a test kit for HCV diagnostics comprising anti-HCV antibodies, HCV virus components and/or cell lines permissive for  
30 HCV replication and expressing these components.

Taken together, the primary object of the present invention is to provide nucleic acids encoding recombinant HCV-derived replicons, which are capable of efficient propagation in a permissive cell culture system.

5 A related object of the invention is to provide susceptible cell lines (and  
"adapted" rapidly growing cell clones derived from them) which support high titer replication  
of recombinant HCV-derived nucleic acids.

By providing cell clones supporting efficient subgenomic HCV replication, the  
present invention provides (i) *in vitro* cell culture models of HCV propagation; (ii) systems  
10 for screening candidate anti-viral compounds and evaluating drug resistance; (iii) methods for  
diagnosing HCV infection, and (iv) systems for production of large quantities of HCV-  
derived components or recombinant viral particles for antibody generation and/or vaccine  
development.

15 **EXAMPLE 1: GENERATION OF STABLE RAPIDLY GROWING HUH-7**  
**HEPATOMA CELL LINES WHICH SUPPORT EFFICIENT**  
**REPLICATION OF SUBGENOMIC HEPATITIS C VIRUS**  
**RNAs**

**Materials and Methods**

*Generation of subgenomic HCV RNAs.* Parental recombinant HCV DNA construct  
20 for expression of subgenomic HCV RNAs was produced by Operon Technologies, Inc.  
(Alameda, CA) using the sequence described by Lohmann *et al.* (Science, 1999, *supra*).

Bicistronic HCV-derived DNA fragment I377/NS3-3'UTR (SEQ ID NO: 1, Figure 3)  
was chemically synthesized and inserted into a modified pUC19 vector under the control of  
T7 promoter to produce expression plasmid pAn/HCVR1. Plasmid DNA was linearized with  
25 *Sca I* and used for *in vitro* transcription reactions with Megascript T7 In Vitro Transcription  
Kit (Ambion) according to manufacturer's instructions. After *in vitro* transcription and  
DNase treatment (to remove template DNA), RNA was extracted with acid phenol  
(Kedzierski and Porter, BioTechniques, 1991, 10:210) and used for transfections.

5       **Cell line growth.** Human hepatoma cell line Huh-7 (Nakabayashi *et al.*, Cancer Res., 1982, 42:3858-3863) was grown in Dulbecco's Modified Eagles Medium (DMEM), supplemented with 10% Fetal Bovine Serum (FBS), Penicillin/Streptomycin, 2 mM glutamine, and non-essential amino acids, in a humidified atmosphere containing 5% CO<sub>2</sub>. Cells were passaged 1-2 times per week at a split ratio of 1:4 – 1:5, using trypsin/EDTA  
10 solution (0.05% / 0.02%). All tissue culture reagents were from Gibco.

**Cell transfection and selection of efficiently growing clones supporting high titer subgenomic HCV replication.** RNA (10-30 µg) was electroporated into  $0.8 \times 10^7$  cells using Gene Pulse instrument (Bio-Rad, 0,4 cm-cuvette, 450V, 960 Fa). Transfection of a construct directing the expression of firefly luciferase was used to optimize transfection efficiency.  
15 Following electroporation, cells were seeded on 10-cm culture dishes in 10 ml of growth medium. After 48 hours, G418 was added at 1 mg/ml, and the medium was changed every second day until (after 3 weeks of selection) resistant colonies were transferred to 96-well plates and passaged 1-2 times a week.

      The fraction of HCV-positive cells in the total population was monitored by  
20 immunofluorescence using polyclonal antibodies raised against non-structural HCV proteins (*see below*) and by RT-PCR (*see below; see also* Lanford *et al.*, Virology, 1994, 202:606; Shimizu *et al.*, Proc. Natl. Acad. Sci. USA, 1992, 89:5477; Mizutani *et al.*, J. Virol., 1996, 70:7219; Ikeda *et al.*, Virus Res., 1998, 56:157; Fournier *et al.*, J. Gen. Virol., 1998, 79:2376).

25       Stable cell clones were expanded and culture conditions were optimized to promote rapid cell growth and HCV RNA replication. Growth rates of the clones were monitored by sulphorhodamine B staining for total cellular protein. The clones which (i) had the growth properties better or indistinguishable from the naive Huh-7 cells (*see, e.g.*, Figure 5) and (ii) supported efficient subgenomic HCV RNA replication (as determined by quantitative RT-  
30 PCR) were chosen for further analysis.

5        **RT-PCR assays.** Total RNA was prepared from transfected cells, and serial dilutions were used for RT-PCR amplification of (i) NS5B region between nt 7435 and 7750 using primers GCCCTAGATTGTCAGATCTACG and ATAAATCCAACCTGGGACGCAGC (SEQ ID NOS: 7 and 8, respectively), (ii) NS5B region between nt 7360 and 7800 using primers CCTTGTGGGCAAGGATGATCC and GACAGGCTGTGATATATGTCTCC (SEQ  
10 ID NOS: 9 and 10, respectively), and (iii) *neo* region between nt 650 and 1110 using primers GTTCTTTTTGTCAAGACCGACC and CCACCATGATATTCGGCAAGC (SEQ ID NOS: 11 and 12, respectively). As a control, RT-PCR was performed without template DNA or using total RNA isolated from naive Huh-7 cells.

15        **Antibodies.** Rabbit polyclonal antisera specific for HCV NS3 or NS5A (Lohmann *et al.*, Science 1999, 285:110-113) were obtained from Ralf Bartenschlager (Institute for Virology, Johannes-Gutenberg University, Mainz, Germany). A mouse monoclonal antibody to HCV NS4B was from Virogen, Inc. (Watertown, MA).

20        **In situ detection of HCV antigens using immunofluorescence.** Monolayer cultures on glass coverslips were fixed with methanol-acetone mixture (1:1) for 2 min, and dried for 10 min at room temperature. Alternatively, cells were fixed directly in 6-well plates (Costar). Dried cell monolayers were blocked with 2% bovine serum albumin (BSA) in phosphate-buffered saline (PBS) for 10 min at room temperature. Double staining of cells was achieved by using rabbit anti-HCV primary antibody followed by the addition of 2 mg/ml of 4',6-diamidino-2-phenylindole (DAPI, Sigma) to stain the cellular DNA, and then secondary  
25 antibody (goat anti-rabbit IgG polyclonal antibody conjugated with rhodamine; Sigma). All primary and secondary antibodies were diluted 1:100 in blocking solution. After mounting, the samples were viewed with an Olympus BX60 microscope with a x20 or x40 objective and specific filter blocks. Images were acquired using a charge-coupled device camera (Olympus, model FKH025144) and processed using Adobe Photoshop and Canvas software.

5        ***Detection of HCV antigens by immunoblotting.*** Monolayer cultures of cells growing on 6-well plates were lysed and subjected to electrophoresis in 12% or 4-20% gradient SDS-polyacrylamide gels (Laemmli, Nature, 1970, 227:680-685). After electrotransfer of proteins to a nitrocellulose filter, HCV antigens were detected using rabbit anti-HCV antibodies and Enhanced Chemiluminescence (ECL) detection kit (Amersham).

10        ***Analysis of subgenomic HCV replicons isolated from rapidly growing Huh-7 cell clones.*** Total RNA was isolated from HCVR 2 (SEQ ID NO: 2), HVCR 8 (SEQ ID NO: 3), HCVR 9 (SEQ ID NO: 4), HCVR 22 (SEQ ID NO: 5) and HCVR 24 (SEQ ID NO: 6 ) rapidly growing Huh-7 cell clones carrying replicating subgenomic HCV constructs. The whole replicon RNA was reverse-transcribed using oligonucleotide primers  
15        CTCGTATGTTGTGTGGAA and GTCGCTCTCGAGGCACATA (SEQ ID NOS: 13 and 14, respectively) designed and synthesized by Operon Technologies (Alameda, CA) and PCR-amplified in overlapping 0.7-1.2 kb-fragments. Total PCR products were sequenced rather than individual cDNA clones, in order to identify predominant mutations. Typically,  
20        every region was sequenced in both directions, using at least two PCR products and several primers for each direction. Sequencing was performed on a contract basis at the Children's Hospital of Philadelphia (Philadelphia, PA). The resulting sequences were analyzed using Sequencher™ 3.1 sequence alignment software.

### **Results and Discussion**

25

Despite the availability of cloned infectious genomes, molecular studies of HCV replication and the development of antiviral drugs have been hampered by the low efficiencies of currently available cell culture systems (*see overview in the Background Section*).

5 To overcome these limitations, an efficient cell culture system was established that is based on the transfection of recombinant HCV-derived RNAs. Using chemical DNA synthesis, pAn/HCVR1 DNA plasmid was constructed, said plasmid encoding a selectable bicistronic replicon (I377/NS3-3'UTR; SEQ ID NO: 1) containing HCV 5' NTR (HCV IRES), the neomycin phosphotransferase (*neo*) gene, the IRES of the encephalomyocarditis virus which directs translation of HCV sequences from NS3 up to NS5B, and HCV 3' NTR (see Materials and Methods and Figure 3).

10 A 5'-flanking T7 RNA polymerase promoter and an engineered restriction site at the 3' end of pAn/HCVR1 allowed for *in vitro* production of run-off RNA transcripts corresponding to a selectable bicistronic replicon. These *in vitro* transcripts (termed "parental" replicons) were transfected into Huh-7 human hepatoma cells. Particular care was taken to remove template DNA, which might otherwise integrate into transfected cells and confer G418 resistance independent of HCV replication (see Materials and Methods). The efficiency of transfection was optimized using luciferase mRNA.

15 In two independent transfection experiments, after 3 weeks of G418 selection (at 1 mg/ml), a total of 33 drug-resistant clones were obtained. These clones were transferred to 96-well plates and passaged 1-2 times a week. Seven cell clones were successfully expanded and were selected for further study.

20 At five weeks posttransfection clones were analyzed by RT-PCR for the presence of HCV RNA (NS5B region was amplified, see Materials and Methods). Clones HCVR 2, 8, 9, 22 and 24 produced PCR fragments of the expected size (Figure 4). These results were confirmed by repeating RT-PCR with three other primer pairs: two corresponding to NS5B, and one to *neo* gene.

25 Although the positive RT-PCR data and the presence of G418 resistance 5-7 weeks posttransfection are strong indications of subgenomic HCV replication, there is still a possibility that both of these effects are due to traces of the input replicon RNA. To rule out this possibility, selected Huh-7 clones which produced positive results by RT-PCR were assayed for the expression of HCV-derived NS proteins. Thus, selected clones HCVR 2, 8, 9, 22 and 24 were analyzed by immunofluorescence microscopy using anti-NS3, NS4, and NS5A antibodies (see Materials and Methods). All three viral proteins were detected in all five cell

5 clones. Similarly to wild-type HCV which replicates in the cytoplasm, the antigens derived from the recombinant replicons of the present invention also localized to the cytoplasm. NS5A protein was expressed in a substantial fraction of cells (at least 30%) in clones HCVR 2 and HCVR 8, but not in naive Huh-7 cells. Similar pattern was observed for NS3, NS4 and NS5A proteins in all five selected cell clones.

10 The presence of HCV replicon-derived antigens in clones HCVR 2, 8 and 9 was also analyzed by Western blotting, and both NS5A and NS3 proteins were detected. Interestingly, when the immunofluorescence detection of NS5A was repeated at 6, 7, 8, 9, and 10 weeks posttransfection, it was found that the fraction of positive cells gradually increased with time, and at 10 weeks posttransfection over 90% of cells (in clones HCVR 8, 9, 22, 24) became  
15 HCV-positive.

Importantly, dramatic changes in cell growth potential were observed over time in several isolated cell clones, which were efficiently replicating subgenomic HCV. Initially (similarly to observations of Lohmann *et al.*, Science, 1999, *supra*), all HCV-replicating cell lines were growing much slower as compared to the parental naive cell line, Huh-7 (*i.e.*,  
20 growth rate of the cell lines was not more than 1% of the growth rate of naive Huh-7 cells). At certain points in time, however, (starting 6 weeks posttransfection) the growth of some of the clones began to accelerate (10-fold or more), ultimately reaching the growth rates of parental Huh-7 cells. As shown in Figure 5, clones HCVR 9 and 24 grew at the same rate as the parental (non-transfected) Huh-7 cells. Similarly, clones HCVR 2, 8 and 22 grew only 4-  
25 fold slower than non-transfected Huh-7 cells. This is in contrast to the growth rates of cells initially prepared by reproducing experiments of Lohmann *et al.*, which had a reduction of growth rate of at least 99% as compared to the naive cell line. Such increase in growth rates of selected clones was not previously reported and coincided with the emergence of detectable HCV-derived antigens and HCV RNA, suggesting that the observed changes were  
30 not associated with the loss of HCV replication.

In the studies of recombinant HCV replicons propagating in Huh-7 cells, Lohmann *et al.* did not observe any time-dependent increase in the efficiency of productive transfection and have not obtained host cell clones with normal growth properties. On the basis of these studies Lohmann *et al.* have hypothesized that HCV RNA replication and/or the expression



5 of HCV NS proteins is always disadvantageous to cells. Moreover, Lohmann *et al.* failed to find any mutations in the 10 replicons that they have sequenced and concluded that formation of an "adapted" replicon would be rare and that the inefficiency of their system is more likely due to particular host cell conditions or factors present in only a few cells.

10 In contrast, the data reported here support a hypothesis that at least some of the selected replicons are "attenuated", *i.e.*, contain mutations that are responsible for their lowered cytotoxicity and high frequency of productive transfection. To identify these mutations, replicon RNAs isolated from all five selected cell clones (HCVR 2, 8, 9, 22 and 24) were amplified using RT-PCR and sequenced. The obtained sequences (SEQ ID NOS: 2-6) were analyzed for the presence of mutations by comparing them to the sequence of the  
15 parental I377/NS3-3'UTR clone (SEQ ID NO: 1). As shown in Table I, mutations are located in various parts of the replicon genome. Further analysis of these mutations may provide important information about the evolution of the virus and the viral genes responsible for replication efficiency and cytotoxicity.

20 **Table I**                      **Mutations in Huh-7 replicon clones**

	Clone#	position	change (nt)	change (aa code)	change aa	Gene
25	HCVR2	1234	C to A			EMCV IRES
	HCVR2	2527	A to G	ATG to GTG	M to V	NS3
	HCVR2	5288	G to T	CGT to CTT	R to L	NS5A
30	HCVR8	2330	A to G	GAA to GGA	E to G	NS3
	HCVR8	after 4847	AAA insertion		added K	NS5A
	HCVR8	5366	C to T	GCA to GTA	A to V	NS5A
35	HCVR9	1236				EMCV IRES
	HCVR9	1905	A to G	CAA to CAG	Silent	NS3
40	HCVR9	3797	A to G	AAG to AGG	K to R	NS4A

5	HCVR9	4364	T to C	GTT to GCT	V to A	NS4B
	HCVR9	4848	C to A	AAC to AAA	N to K	NS5A
10	HCVR22	2330	A to G	GAA to GGA	E to G	NS3
	HCVR22	3935	A to G	CAA to CGA	Q to R	NS4B
	HCVR22	5320	G to A	GCC to ACC	A to T	NS5A
15	HCVR24	1760	C to T			EMCV RES
	HCVR24	3016	G to T	GAC to TAC	D to Y	NS3
20	HCVR24	5336	G to T	AGC to ATC	S to I	NS5A

In summary, in contrast to all previous reports, using prolonged selection in the presence of G418, the present inventors have generated HCV-replicating "adapted" cell clones, which grow at the rate, which is similar to or indistinguishable from the non-transfected parental Huh-7 cell line. These cell clones support efficient propagation of recombinant HCV replicon mutants at least some of which have unusually low cytotoxicity leading to high replication titers and high cell growth rate.

Cell clones HCVR 2, 8, 9, 22 and 24, were deposited with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209, USA on September 20, 2000, and assigned Accession Nos. as set forth below:

ATCC Deposits	
HCV cell clone	Accession No.
HCVR 2	PTA-2489
HCVR 8	PTA-2490
HCVR 9	PTA-2486
HCVR 22	PTA-2487
HCVR 24	PTA-2488

5

**EXAMPLE 2:      SCREENING ASSAYS FOR ANTI-HCV THERAPEUTICS**  
**USING HCV-REPLICATING CELL CLONES OF**  
**THE PRESENT INVENTION**

10

In the identified efficiently growing "adapted" Huh-7 cell clones containing replicating recombinant HCV genomes, conditions are optimized for cell growth and HCV RNA replication. The timing of exposure to test compounds is determined based on the kinetics of HCV RNA accumulation and/or decline in the absence of the selective agent, and  
15 the effects of various known inhibitors of RNA synthesis. A cell-based assay for HCV genome replication is developed based on all these data. In this assay, candidate compounds are tested for inhibition of HCV RNA replication that is selective with respect to inhibition of cell growth. In particular, shifting of growth rate curves to a lower growth rate in the presence of a test compound indicates that the compound is a potential lead for developing an  
20 anti-HCV therapeutic.

\* \* \* \* \*

25 The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and the accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

30 All values given in the specification are approximate, and are provided for illustration and not by way of limitation.

5 **SEQUENCE LISTING**

## 1. Parental subgenomic HCV replicon I377/NS3-3'UTR

SEQ ID NO: 1

1 gccagccccc gatgggggc gacactccac catagatcac tccctgtga ggaactactg  
 61 tcttcacgca gaaagcgtct agccatggcg ttagtatgag tgcgtgcag cctccaggac  
 10 121 ccccccctccc gggagagcca tagtggctcg cggaaccggt gagtacaccg gaattgccag  
 181 gacgaccggg tctttcttg gatcaaccg ctcaatgcct ggagatttgg gcgtgcccc  
 241 gcgagactgc tagccgagta gtgtgggtc gcgaaaggcc ttgtgtact gcctgatagg  
 301 gtgcttgcga gtccccggg aggtctcgt gaccgtgcac catgagcacg aatcctaaac  
 361 ctcaaagaaa aaccaaaggg cgcgccatga ttgaacaaga tggattgcac gcaggttctc  
 15 421 cggccgcttg ggtggagagg ctattcggct atgactgggc acaacagaca atcggctgct  
 481 ctgatccgc cgtgttccg ctgtcagcg aggggcgccc ggttctttt gtcaagaccg  
 541 acctgtccg tgcctgaat gaactgcagg acgaggcagc gcggctatcg tggctggcca  
 601 cgacggcgct tcttgcgca gctgtgctcg acgtgtcac tgaagcgga agggactggc  
 661 tgctattggg cgaagtgcg gggcaggatc tctgtcatc tcacctgct cctgccgaga  
 20 721 aagtatccat catggctgat gcaatgcggc ggctgcatac gcttgatccg gctacctgcc  
 781 cattogacca ccaagcgaac catgcacgc agcgagcacg tactcggatg gaagccggtc  
 841 ttgtcatca ggatgatcg gacgaagagc atcaggggct cgcgccagcc gaactgttcg  
 901 ccaggctcaa ggcgcgatg cccgacggcg aggatctcgt cgtgacctat ggcatgcct  
 961 gcttgccgaa tatcatggtg gaaaatggc gctttcttg attcatcgac tgtggccggc  
 25 1021 tgggtgtggc ggaccgctat caggacatag cgttggctac ccgtgatatt gctgaagagc  
 1081 ttggcggcga atgggtgac cgcttctcgc tgccttaccg tatcggcgt cccgattcgc  
 1141 agcgcatcgc ctctatcgc ctcttgacg agttctctg agtttaaaca gaccacaacg  
 1201 gttccctct agcgggatca attccgccc tctccctccc cccccctaa cgttactggc  
 1261 cgaagccgct tggaataagg ccggtgtgcg ttgtctata tgtatttcc caccatattg  
 30 1321 ccgtcttttg gcaatgtgag ggcccggaaa cctggccctg tcttctgac gagcattcct  
 1381 aggggtcttt cccctctgc caaaggaatg caaggtctgt tgaatgtcgt gaaggaagca  
 1441 gttctcttg aagcttctg aagacaaca acgtctgtg cgacctttg caggcagcgg  
 1501 aacccccac ctggcgacag gtgcctctgc ggccaaaagc cacgtgtata agatacacct  
 1561 gcaaaggcgg cacaaccca gtgccacgtt gtgagtggga tagttgtgga aagagtcaaa

5 1621 ttgctctcct caagcgtatt caacaagggg ctgaaggatg cccagaaggt accccattgt  
 1681 atgggatctg atctggggcc tcggtgcaca tgctttacat gtgtttagtc gaggttaaaa  
 1741 aacgtctagg cccccgaac cacggggacg tggttttcct ttgaaaaaca cgataatacc  
 1801 atggcgccta ttacggccta ctcccaacag acgcgaggcc tacttggctg catcatcact  
 1861 agcctcacag gccgggacag gaaccaggctc gagggggagg tccaagtgtt ctccaccgca  
 10 1921 acacaatctt tcctggcgac ctgcgtcaat ggcgtgtgtt ggactgtcta tcatggtgcc  
 1981 ggtcacaaga cccttgccgg cccaaagggc ccaatcacc aaatgtacac caatgtggac  
 2041 caggacctcg tcggctggca agcggcccc ggggcgcgtt ccttgacacc atgcacctgc  
 2101 ggcagctcgg acctttactt ggtcacgagg catgccgatg tcattccggt gcgccggcgg  
 2161 ggcgacagca gggggagcct actctcccc aggcccgctt cctacttgaa gggctcttcg  
 15 2221 ggcgtccac tgctctgcc ctggggcac gctgtgggca tcttcgggc tgccgtgtgc  
 2281 acccgagggg ttgcgaaggc ggtggacttt gtaccgctg agtctatgga aaccactatg  
 2341 cggtccccgg tcttcacgga caactcgtcc cctccggccg tacgcagac attccaggtg  
 2401 gccatctac acgccctac tggtagcggc aagagcacta aggtgccggc tgcgtatgca  
 2461 gcccaagggt ataaggtgct tgcctgaac ccgtccgtg ccgccaccct aggtttcggg  
 20 2521 gcgtatatgt ctaaggcaca tggatcgac cctaacaatca gaaccggggt aaggaccatc  
 2581 accacgggtg ccccatcac gtactccacc tatggcaagt ttcttgcca cggtggttgc  
 2641 tctggggcg cctatgacat cataatatgt gatgagtcc actcaactga ctgcaccact  
 2701 atctgggca tcggcacagt cctggaccaa gcggagacgg ctggagcgcg actcgtcgtg  
 2761 ctgccaccg ctacgcctcc gggatcggtc accgtgccac atccaaacat cgaggagggtg  
 25 2821 gctctgtcca gcaactggaga aatccccctt tatggcaaag ccatcccat cgagaccatc  
 2881 aaggggggga ggcacctcat ttctgcat tccaagaaga aatgtgatga gctcgccgcg  
 2941 aagctgtccg gcctcggact caatgtgta gcatattacc ggggccttga tgtatccgtc  
 3001 ataccaacta gcggagacgt cattgtcgt gcaacggacg ctctaatac gggctttacc  
 3061 ggcatctcg actcagtgt cgactgcaat acatgtgtca cccagacagt cgacttcagc  
 30 3121 ctggaccga ccttaccat tgagacgac accgtgccac aagacgcgtt gtcacgctc  
 3181 cagcggcgag gcaggactgg taggggcagg atgggcattt acaggtttgt gactccagga  
 3241 gaacggccct cgggcatgtt cgattcctc gtctgtgcg agtgctatga cgcgggctgt  
 3301 gcttggtac agctcacgcc cgccgagacc tcagttaggt tgcgggctta cctaaacaca  
 3361 ccagggttgc ccgtctgcca ggaccatctg gagtctggg agagcgtctt tacaggcctc

5 3421 acccacatag acgcccattt ctgtcccag actaagcagg caggagacaa ctccccctac  
 3481 ctggtagcat accaggctac ggtgtgcgc agggctcagg ctccacctcc atcgtgggac  
 3541 caaatgtgga agtgtctcat acggctaaag cctacgtgc acgggccaac gcccctgctg  
 3601 tataggctgg gagccgttca aaacgaggtt actaccacac accccataac caatacatc  
 3661 atggcatgca tgtcggctga cctggaggtc gtcacgagca cctgggtgct ggtaggcgga  
 10 3721 gtcctagcag ctctggccgc gtattgcctg acaacaggca gcgtggtcat tgtgggcagg  
 3781 atcatctgt ccggaagcc ggccatcatt ccgacaggg aagtcctta ccgggagttc  
 3841 gatgagatgg aagagtgcgc ctcacacctc cctacatcg aacagggaat gcagctcgcc  
 3901 gaacaattca aacagaaggc aatcgggttg ctgcaaacag ccaccaagca agcggaggct  
 3961 gctgctccc tggtggaatc caagtggcgg accctgaag ccttctgggc gaagcatatg  
 15 4021 tggaaattca tcagcgggat acaatattta gcaggcttgt ccactctgcc tggcaacccc  
 4081 gcgatagcat cactgatggc attcacagcc tctatcacca gcccgctcac cacccaacat  
 4141 accctctgt ttaacatcct ggggggatgg gtggcgcgc aacttgctcc tccagcgct  
 4201 gcttctgct tcgtaggcgc cggtatcgt ggagcggctg ttggcagcat aggccttggg  
 4261 aagtgcttg ttgatattt ggaggttat ggagcagggg ttgcagcgc gctcgtggcc  
 20 4321 ttaaggtca tgagcggcga gatgccctcc accgaggacc tggtaacct actcctgct  
 4381 atctctccc ctggcgccct agtcgtcggg gtcgtgtcg cagcgatact gcgtcggcac  
 4441 gtggcccag gggagggggc tgtcagtgg atgaaccggc ttagagcgtt cgcttcggc  
 4501 ggtaaccacg tctccccac gcactatgtg cctgagagcg acgtgcagc acgtgtcact  
 4561 cagatcctct ctagtctac catcactcag ctgtgaaga ggcttcacca gtggatcaac  
 25 4621 gaggactgct ccacgccatg ctccggctcg tggtaagag atgtttgga ttgatatgc  
 4681 acggtgttga ctgattcaa gacctggctc cagtccaagc tctgccgcg attgccggga  
 4741 gtccccctct tctcatgca acgtgggtac aaggagctt ggccgggcga cggcatcatg  
 4801 caaacacct gccatgtgg agcacagatc accggacatg tgaaaaacgg ttccatgagg  
 4861 atcgtggggc ctaggacctg tagtaacacg tggcatgga cattcccat taacgcgtac  
 30 4921 accacgggcc cctgcacgcc ctccccggcg ccaaattatt ctaggcgct gtggcgggtg  
 4981 gctgctgagg agtacgtgga ggttacggg gtgggggatt tccactacgt gacgggcatg  
 5041 accactgaca acgtaaagt cccgtgtcag gtccggccc ccgaattctt cacagaagtg  
 5101 gatggggtgc ggtgcacag gtacgtcca gcgtgcaaac cctcctacg ggaggaggtc  
 5161 acattcctg tcgggctcaa tcaatacctg gtgggtcac agctcccatg cgagcccgaa

5 5221 ccggacgtag cagtgtcac ttccatgctc accgaccctt cccacattac ggaggagacg  
 5281 gctaagcgta ggctggccag gggatctccc cctccttgg ccagctcatc agctagccag  
 5341 ctgtctgcgc ctctctgaa ggcaacatgc actaccgtc atgactcccc ggacgtgac  
 5401 ctatcgagg ccaacctct gtggcggcag gagatggcg ggaacatcac ccgctggag  
 5461 tcagaaaata aggtagtaat ttggactct ttcgagccgc tccaagcgga ggaggatgag  
 10 5521 agggaaagtat ccgtccggc ggagatcctg cggaggtcca ggaaattccc tcgagcgtg  
 5581 cccatattggg cagcccgga ttacaacct cactgttag agtctggaa ggaccggac  
 5641 tacgtccctc cagtgttaca cgggtgtcca ttccgcctg ccaagcccc tccgatacca  
 5701 cctccacgga ggaaggagc ggtgtcctg tcagaatcta ccgtgtctt tccttggcg  
 5761 gagctcgcca caaagacct cggcagctcc gaatcgtcg ccgtcgacag cggcacggca  
 15 5821 acggcctctc ctgaccagcc ctccgacgac ggcgacggg gatccgacgt tgagtgtac  
 5881 tctccatgc cccccctga gggggagccg ggggatccg atctcagca cgggtcttg  
 5941 tctaccgtaa gcgaggaggc tagtgaggac gtcgtctgt gtcgatgtc ctacacatgg  
 6001 acaggcgccc tgatcacgc atgcgtgcg gaggaacca agctgccat caatgactg  
 6061 agcaactct tgctccgta ccacaactg gtctatgcta caacatctg cagcgcaagc  
 20 6121 ctgcggcaga agaaggtcac cttgacaga ctgcaggtcc tggacgacca ctaccgggac  
 6181 gtgtcaagg agatgaaggc gaaggctcc acagttaagg ctaaactct atccgtggag  
 6241 gaagcctgta agctgacgc cccacattc gccagatcta aatttgcta tggggcaaag  
 6301 gacgtccgga acctatccag caaggccgtt aaccacatc gctccgtgt gaaggactg  
 6361 ctggaagaca ctgagacacc aattgacacc accatcatg caaaaatga ggtttctgc  
 25 6421 gtccaaccag agaaggggg ccgcaagcca gtcgcctta tctattccc agattgggg  
 6481 gttcgtgtg gcgagaaaat ggcccttac gatgtggtt ccacctccc tcaggccgtg  
 6541 atgggtctt catagcgtt ccaatactt cctggacgc ggtcgagt cctgtgaat  
 6601 gcctggaaag cgaagaaat ccctatggc ttgcatac acaccgctg tttgactca  
 6661 acggtcactg agaatgacat ccgtgtgag gactcaatc accaatgtt tgactggcc  
 30 6721 cccgaagcca gacaggccat aaggtcgtc acagagcggc ttacatcgg gggccccctg  
 6781 actaattcta aagggcagaa ctgcggctat cggcggtgcc gcgcgagcgg tgtactgac  
 6841 accagctgcg gtaataacct cacatgttac ttgaaggccg ctgcggcctg tcgagctgcg  
 6901 aagctccagg actgcacgt gctcgtatg ggagacgacc ttgtgttat ctgtgaaagc  
 6961 gcggggaccc aaggaggca ggcgagccta cgggcctta cggaggctat gactagatac

5 7021 tctcccccc ctggggaccc gcccaaacca gaatacgact tggagtgtat aacatcatgc  
 7081 tcctccaatg tgtcagtcgc gcacgatgca tctggcaaaa ggggtgtacta tctcaccgt  
 7141 gacccacca ccccccttgc gcgggctgcg tgggagacag ctagacacac tccagtcaat  
 7201 tcctggctag gcaacatcat catgtatgcg cccacctgtt gggcaaggat gatcctgaig  
 7261 actcatttct tctccatcct tctagctcag gaacaacttg aaaaagccct agattgtcag  
 10 7321 atctacgggg cctgttactc cattgagcca ctgacctac ctcagatcat tcaacgactc  
 7381 catggcctta gcgcatttc actccatagt tactctccag gtgagatcaa taggggtggct  
 7441 tcatgcctca ggaaacttgg ggtaccgccc ttgcgagtct ggagacatcg ggccagaagt  
 7501 gtccgcgcta ggctactgtc ccaggggggg agggctgcca ctgtggcaa gtacctcttc  
 7561 aactgggcag taaggaccaa gctcaaactc actccaatcc cggctgcgtc ccagttggat  
 15 7621 ttatccagct ggttcgttgc tggttacagc gggggagaca tatatcacag cctgtctcgt  
 7681 gcccgacccc gctggttcat gtggtgccta ctctacttt ctgtaggggt aggcatttat  
 7741 ctactcccca accgatgaac ggggagctaa aactccagg ccaataggcc atcctgtttt  
 7801 ttccctttt tttttctt tttttttt tttttttt tttttttt tttctcttt  
 7861 ttttctct tttttctt tttttctt tgggtgctcc atcttagccc tagtcacggc  
 20 7921 tagctgtgaa aggtccgtga gccgcttgac tgcagagagt gctgatactg gcctctctgc  
 7981 agatcaagta ct

LOCUS SSE242652 7989 bp RNA SYN 07-JUL-1999

DEFINITION: Hepatitis C virus replicon I377/NS3-3'UTR.

25 ACCESSION: AJ242652

NID g5441834

VERSION AJ242652.1 GI:5441834

30 KEYWORDS: core-neo gene; NS2 gene; NS2 proteinase; NS3 gene; NS3  
 proteinase/helicase; NS3/4A proteinase cofactor; NS4A gene; NS4B gene; NS4b protein; NS5a  
 gene; NS5A phosphoprotein; NS5b gene; NS5B RNA dependant RNA polymerase; polyprotein.

SOURCE: Hepatitis C virus replicon I377/NS3-3'UTR.

ORGANISM: Hepatitis C virus replicon I377/NS3-3'UTR artificial sequence; vectors.

REFERENCE: 1 (bases 1 to 7989)



5   AUTHORS:       Lohmann,V.,   Korner,F.,   Koch,J.,   Herian,U.,   Theilmann,L.   and  
      Bartenschlager,R.  
      TITLE:        Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line  
      JOURNAL:       Science 285 (5424), 110-113 (1999)  
      MEDLINE        99322193

10   REFERENCE:     2 (bases 1 to 7989)  
      AUTHORS:       Bartenschlager,R.  
      TITLE:        Direct Submission  
      JOURNAL:       Submitted (26-MAY-1999) Bartenschlager R., Institute for Virology,  
      Johannes Gutenberg - University Mainz, Obere Zahlbacher Strasse 67, 55131 Mainz, GERMANY

15   FEATURES:       Location/Qualifiers  
      SOURCE:        1..7989  
                     /organism="Hepatitis C virus replicon I377/NS3-3'UTR"  
                     /note="Neomycin-selectable bicistronic subgenomic  
                     hepatitis C virus replicon"  
      20             /db\_xref="taxon:95363"  
          source     1..7989  
                     /organism="Encephalomyocarditis virus"  
                     /db\_xref="taxon:12104"  
      5'UTR         1..341  
      25             /evidence=experimental  
          source     1..7989  
                     /organism="Hepatitis C virus"  
                     /db\_xref="taxon:11103"  
          source     join(1..341,1801..7758)  
      30             /organism="Hepatitis C virus"  
                     /strain="1b"  
                     /isolate="Con1"  
                     /db\_xref="taxon:11103"  
                     /tissue\_type="liver"

5       gene       342..1181  
          /feature="core-neo"  
      CDS       342..1181  
          /feature="core-neo"  
          /function="selectable marker"  
10       /codon\_start=1  
          /product="core-neo fusion protein"  
          /protein\_id="CAB46912.1"  
          /db\_xref="PID:e1512562"  
          /db\_xref="PID:g5441835"  
15       /db\_xref="GI:5441835"

/translation="MSTNPKPQRKTKGRAMIEQDGLHAGSPAAWVERLFGYDWAQQTIGCS  
DAAVFRLSAQGRPVLFVKTDLSGALNELQDEAARLSWLATTGVPCAAVLDVVTEAG  
RDWLLLGEVPGQDLLSSHLAPAELKVSIMADAMRRLHTLDPATCFDHQAKHRIERA  
20 RTRMEAGLVDQDDLDEEHQGLAPAELFARLKARMPDGEDLVVTHGDACLPNIMVE  
NGRFSGFIDCGRLGVADRYQDIALATRDIAEELGGEWADRFLVLYGIAAPDSQRIAFY  
RLLDEFF"

      misc\_feature 1190..1800  
          /feature="internal ribosome entry site (IRES)"  
25       source       1190..1800  
          /organism="Encephalomyocarditis virus"  
          /db\_xref="taxon:12104"  
      gene       1801..3696  
          /feature="NS3"  
30       mat\_peptide 1801..3696  
          /feature="NS3"  
          /product="NS3 proteinase/helicase"  
      CDS       1801..7758  
          /codon\_start=1

5           /product="non-structural polyprotein"  
          /protein\_id="CAB46913.1"  
          /db\_xref="PID:e1512563"  
          /db\_xref="PID:g5441836"  
          /db\_xref="GI:5441836"

10           /translation="MAPITAYSQQTRGLLGCIITSLTGRDRNQVEGEVQVVSTATQSFLATCV  
NGVCWTVYHGAGSKTLAGPKGPITQMYTNVDQDLVGWQAPPGARSLTPCTCGSSD  
LYLVTRHADVIPVRRRGDSRGSLLSPRPVSYLKGSSGGPLLCPSGHAVGIFRAAVCTR  
GVAKAVDFVPVESMETTMRSPVFTDNSSPPAVPQTFQVAHLHAPTGS GKSTKVPAA  
15 YAAQGYKVLVLNPSVAATLGFGAYMSKAHGIDPNIRTGVRTTTTGAPITYSTYGKFL  
ADGGCSGGAYDIICDECHSTDSTTILGIGTVLDQAETAGARLVVLATATPPGSVTV  
HPNIEVALSSTGEIPFYGKAIPETIKGGRHLIFCHSKKKCDELAAKLSGLGLNAVAY  
YRGLDVSVIPTSGDVIVVATDALMTGFTGDFDSVIDCNTCVTQTVDFSLDPTFTIETTT  
VPQDAVSRSQRRGRTGRGRMGIIYRFVTPGERPSGMFSSVLCECYDAGCAWYELTP  
20 AETSVRLRAYLNTPLPVCQDHLEFWESVFTGLTHIDAHFLSQTQAGDNFPYLVA  
QATVCARAQAPPPSWDQMWKCLIRLKPTLHGPTPLLYRLGAVQNEVTTHHPITKYIM  
ACMSADLEVVTSTWVLVGGVLAALAAAYCLTTGSVVIVGRIILSGKPAIIPDREVL  
YRE FDEMEECASHLPYIEQGMQLAEQFKQKAIGLLQTATKQAEAAAPVVESKWRTLEAF  
WAKHMWNFISGIQYLAGLSTLPGNPAIASLMAFTASITSPLTTQHTLLFNILGGWVAA  
25 QLAPPSAASAFVGAGIAGAAVGSIGLGKVLVDILAGYGAGVAGALVAFKVMSEMP  
STEDLVNLLPAILSPGALVVGVCVCAAILRRHVGPGEVAVQWMNRLIAFASRGNHVSP  
THYVPESDAAARVTQILSSLTITQLLRLHQWINE DCSTPCSGSWLRD VWDWICTVL  
TDFKTWLQSKLLPRLPGVPPFFSCQRGYKGVWRGDGIMQTTCPGAQITGHVKNGSM  
RIVGPRTCSNTWHGTFPINAYTTGPCTPSPAPNYSRALWRVAAEEYVEVTRVGD  
30 VTGMTTDNVKCPQVPAPEFFTEVDGVRLHRYAPACKPLLREEVTFLVGLNQYLV  
SQLPCEPEPDVAVLTSMLTDP SHITAETA KRRLARGSPPSLASSASQLSAPSLKATCT  
TRHDSPADLIEANLLWRQEMGGNITRVESENKVVILDSFEPLQAEEDEREVSVP  
AEI LRRSRKFPRAMPIWARPDYNPPLLESWKDPDYVPPVVHGCPLPPAKAPP  
IPPPRRKRT VVLSESTVSSALAEATKTFGSSESAVDSGTATASPDQPSDDGDAGSDVESYSSMPP

5 LEGEPGDPDLSDGSWSTVSEEBASEDVVCCSMSYTWGTALITPCAAEETKLPINALSNS  
LLRHHNLVYATTSRSASLRQKKVTFDRLQVLDDHYRDVLKEMKAKASTVKAKLLS  
VEEACKLTPPHSARSKFGYGAKDVRLSSKAVNHRSVWKDLLEDTPIDTTIMAK  
NEVFCVQPEKGGRKPARLIVFIDLGVVRVCEKMALYDVVSTLPQAVMGSSYGQYSP  
GQRVEFLVNAWKAKKCPMGFAYDTRCFDSTVTENDIRVEESIQCCDLAPEARQAIR  
10 SLTERLYIGGPLTNSKGQNCGYRRCRASGVLTTSCGNTLTCYLKAAAACRAAKLQD  
CTMLVCGDDLVICESAGTQEDEASLRAFTEAMTRYSAAPPDPPKPEYDLELITSCSS  
NVSVAHDASGKRVYYLTRDPTTPLARAAWETARHTPVNSWLGNIMYAPTLWARM  
LMTHFFSILLAQEQLEKALDCQIYGACYSIEPLDLPQIIQRLHGLSAFSLHSYSPGEINR  
VASCLRKLGVPLRVWRHRARSVRARLLSQGGRAATCGKYLFWAVRTKLKLTPIP  
15 AASQLDLSSWFVAGYSGGDIYHLSRARPRWFMWCLLLLSVGVGIYLLPNR"

gene 3697..3858  
/gene="NS4A"  
mat\_peptide 3697..3858  
20 /gene="NS4A"  
/product="NS3/4A proteinase cofactor"  
mat\_peptide 3859..4641  
/gene="NS4B"  
/product="NS4B protein"  
25 gene 3859..4641  
/gene="NS4B"  
mat\_peptide 4642..5982  
/gene="NS5A"  
/product="NS5A phosphoprotein"  
30 gene 4642..5982  
/gene="NS5A"  
mat\_peptide 5983..7755  
/gene="NS5B"  
/product="NS5B RNA dependant RNA polymerase"

5 gene 5983..7755  
/gene="NS5B"

3'UTR 7759..7989

BASE COUNT 1647 a 2368 c 2243 g 1731 t

ORIGIN

10

## 2. HCV Replicon RNA from cell line HCVR2

SEQ ID NO: 2

1 gccagccccc gattgggggc gacactccac catagatcac tcccctgtga ggaactactg  
15 61 ttctcacgca gaaagcgtct agccatggcg ttagtatgag tgcgtgcag cctccaggac  
121 cccccctccc gggagagcca tagtggtctg cggaaccggt gactacaccg gaattgccag  
181 gacgaccggg tcctttcttg gatcaaccgc ctcaatgcct ggagattgg gcgtgcccc  
241 gcgagactgc tagccgagta gtgtgggtc gcgaaaggcc ttgtgtact gcctgatagg  
301 gtgcttgcga gtgccccggg aggtctcgta gaccgtgcac catgagcacg aatcctaaac  
20 361 ctcaaagaaa aaccaaaggc gcgcctatga ttgaacaaga tggattgcac gcaggttctc  
421 cggccgcttg ggtggagagg ctattcggct atgactgggc acaacagaca atcggtgct  
481 ctgatccgc cgtgtccgg ctgtcagcg aggggcgccc ggttctttt gtcaagaccg  
541 acctgtccgg tgccctgaat gaactgcagg acgaggcagc gcggctatcg tggctggcca  
601 cgacgggcgt tccttgcgca gctgtgctcg acgtgtcac tgaagcggga agggactggc  
25 661 tgctattggg cgaagtgcg gggcaggatc tcctgtcatc tcacctgct cctgccgaga  
721 aagtatccat catggctgat gcaatcggc ggctgcatac gctgatccg gctacctgcc  
781 cattcgacca ccaagcgaac catcgcatc agcgagcacg tactcggatg gaagccggtc  
841 ttgtgatca ggatgatctg gacgaagagc atcaggggct cgcgccagcc gaactgttcg  
901 ccaggctcaa ggcgcgatg cccgacggcg aggatctcgt cgtgacccat ggcgatgcct  
30 961 gcttgccgaa tatcatgggt gaaaatggcc gctttctgg attcatcgac tgtggccggc  
1021 tgggtgtggc ggaccgctat caggacatag cgttggctac cgtgatatt gctgaagagc  
1081 ttggcggcga atgggctgac cgttcctcg tgctttacgg tatcggcgt cccgattcgc  
1141 agcgcacgc ctctatcgc ctcttgacg agttctctg agtttaaca gaccacaacg  
1201 gtttcctct agcgggatca attccgccc tctAacctccc cccccctaa cgttactggc

5 1261 cgaagccgct tggaataagg ccggtgtgcg ttgtctata tgtattttc caccatattg  
 1321 ccgtcttttg gcaatgtgag ggcccggaaa cctggccctg tcttctgac gagcattcct  
 1381 aggggtcttt cccctctgc caaaggaatg caaggtctgt tgaatgtcgt gaaggaagca  
 1441 gtctctctgg aagcttcttg aagacaaaca acgtctgtag cgacccttg caggcagcgg  
 1501 aacccccac ctggcgacag gtgcctctgc ggccaaaagc cacgtgtata agatacacct  
 10 1561 gcaaaggcgg cacaacccca gtgccacgtt gtgagtggga tagttgtgga aagagtcaaa  
 1621 ttggtctcct caagcgtatt caacaagggg ctgaaggatg ccagaagggt accccattgt  
 1681 atgggatctg atctggggcc tcggtgcaca tgccttacct gtgttagtc gaggttaaaa  
 1741 aacgtctagg ccccccgaac cacggggacg tggtttctt ttgaaaaaca cgataatacc  
 1801 atggcgccca ttacggccca ctccaacag acgcgaggcc tacttggctg catcatcact  
 15 1861 agcctcacag gccgggacag gaaccaggtc gagggggagg tccaagtgtt ctccaccgca  
 1921 acacaatctt tcttggcgac ctgcgtcaat ggcgtgtgtt ggactgtcta tcatggtgcc  
 1981 ggctcaaaaga ccttggccgg ccaaagggc ccaatcacc aaatgtacac caatgtggac  
 2041 caggacctcg tcggctggca agcgcacccc ggggcgcgtt ccttgacacc atgcacctgc  
 2101 ggcagctcgg accttacct ggtcacgagg catgccgatg tcatccggt gcgccggcgg  
 20 2161 ggcgacagca gggggagcct actctcccc aggcccgctt cctactgaa gggctcttcg  
 2221 ggcgtccac tgctctgccc ctggggcac gctgtgggca tcttcgggc tgcgtgtgc  
 2281 acccgagggg ttgcgaaggc ggtggacttt gtaccgctg agtctatgga aaccactatg  
 2341 cggccccggg tcttcacgga caactcgtcc cctccggccg taccgcagac attccagggtg  
 2401 gcccattac acgcccctac tggtagcggc aagagcacta aggtgccggc tgcgtatgca  
 25 2461 gccaagggt ataagggtgt tgcctgaac ccgtccgtc cgcaccctt aggtttcggg  
 2521 gcgtatGtgt ctaaggcaca tggatcgac ctaaatca gaaccggggt aaggaccatc  
 2581 accacgggtg ccccatcac gtactccacc tatggcaagt ttctgccga cgggtgttgc  
 2641 tctgggggag cctatgacat cataatatgt gatgagtgc actcaactga ctgcaccact  
 2701 atctgggca tcggcacagt cctggaccaa gcggagacgg ctggagcgcg actcgtcgtg  
 30 2761 ctgccaccg ctacgcctcc gggatcggtc accgtgccac atccaaacat cgaggagggtg  
 2821 gctctgtcca gactggaga aatccccctt tatggcaaag ccatcccat cgagaccatc  
 2881 aaggggggga ggcacctcat ttctgccaat tccaagaaga aatgtgatga gctcggcgcg  
 2941 aagctgtccg gcctcggact caatgctgta gcatattacc ggggccttga tgtatccgtc  
 3001 ataccaacta gcggagacgt cattgtcgta gcaacggacg ctctaatac gggctttacc

5     3061 ggcgatttcg actcagtgat cgactgcaat acatgtgtca cccagacagt cgacttcagc  
       3121 ctggaccgga ccttcacat tgagacgacg accgtgccac aagacgcggt gtcacgctcg  
       3181 cagcggcgag gcaggactgg taggggcagg atgggcattt acaggtttgt gactccagga  
       3241 gaacggccct cgggcatgtt cgattcctcg gttctgtgcg agtgctatga cgcgggctgt  
       3301 gcttggtacg agctcacgcc cgccgagacc tcagttaggt tgcgggctta cctaaacaca  
 10     3361 ccagggttgc cgtctgcca ggaccatctg gattctggg agagcgtctt tacaggcctc  
       3421 accacatag acgccattt ctgtcccag actaagcagg caggagacaa ctccccctac  
       3481 ctggtagcat accaggctac ggtgtgcgcc agggctcagg ctccacctcc atcgtgggac  
       3541 caaatgtgga agtgtctcat acggctaaag cctacgtgc acgggccaac gccctgtcg  
       3601 tataggctgg gagccgttca aaacgaggtt actaccacac acccataac caatacatc  
 15     3661 atggcatgca tgtcggctga cctggaggtc gtcacgagca cctgggtgct ggtaggcgga  
       3721 gtcttagcag ctctggccgc gtattgcctg acaacaggca gcgtggcat tgtgggcagg  
       3781 atcatctgt cgggaaagcc ggccatcatt ccgacaggg aagtcctta cggggagtic  
       3841 gatgagatgg aagagtgcgc ctacacctc cttacatcg aacagggaat gcagctcgcc  
       3901 gaacaattca aacagaaggc aatcgggttg ctgaaacag ccaccaagca agcggaggct  
 20     3961 gctgctcccg tggtggaatc caagtggcgg accctgaag cttctgggc gaagcatatg  
       4021 tggaaattca tcagcgggat acaatattha gcaggctgt cactctgcc tggcaacccc  
       4081 gcgatacat cactgatggc attcacagcc tctatacca gcccgctac cacccaacat  
       4141 accctcctgt ttaacatct ggggggatgg gtggccgcc aacttgctc tccagcgt  
       4201 gcttctgctt tcgtaggcgc cggcatcgct ggagcggctg ttggcagcat aggccttggg  
 25     4261 aagggtgctg tggatattht ggaggttat ggagcagggg tggcaggcgc gctcgtggcc  
       4321 ttaaggta tgagcggcga gatgccctcc accgaggacc tggtaacct actccctgct  
       4381 atcctctccc ctggcgcct agtcgtcggg gtcgtgtgcg cagcgatact gcgtcggcac  
       4441 gtgggcccag gggagggggc tgtcagtggt atgaaccggc tgatagcgtt cgcttcgcg  
       4501 ggtaaccacg tctccccac gcactatgt cctgagagcg acgtgcagc acgtgtcact  
 30     4561 cagatcctct ctagtcttac catcactcag ctgctgaaga ggcttcacca gtggatcaac  
       4621 gaggactgct ccacgccatg ctccggctcg tggctaagag atgttggga ttggatatgc  
       4681 acgggtgtga ctgattcaa gacctggctc cagtccaagc tctgccgcg attgccggga  
       4741 gtccctctt tctcatgta acgtgggtac aaggaggtct ggccgggcga cggcatcatg  
       4801 caaaccacct gcccatgtgg agcacagatc accggacatg tgaaaacgg ttcatgagg

5 4861 atcgtggggc ctaggacctg tagtaacacg tggcatggaa cattcccat taacgcgtac  
 4921 accacgggcc cctgcacgcc ctcccggcg ccaaattatt ctaggcgct gtggcgggtg  
 4981 gctgctgagg agtacgtgga ggttacgcgg gtgggggatt tccactacgt gacgggcatg  
 5041 accactgaca acgtaaagtg cccgtgtcag gtccggccc ccgaattctt cacagaagtg  
 5101 gatgggggtg ggttcacag gtacgtcca gcgtgcaa cctcctacg ggaggaggtc  
 10 5161 acattcctgg tcgggctcaa tcaatactg gttgggtcac agtcccatg cgagcccgaa  
 5221 ccggacgtag cagtgtcac ttcatgctc accgaccctt ccacattac ggaggagacg  
 5281 gtaagcTta gggtggccag gggatctccc cctccttgg ccagctatc agtagccag  
 5341 ctgtctgcgc cticctgaa ggcaacatgc actaccgtc atgactccc ggacgtgac  
 5401 ctcatgagg ccaacctct gtggcgagc gagatggcg ggaacatcac ccgctggag  
 15 5461 tcagaaaata aggtagtaatt ttggactct ttcagccgc tccaagcgga ggaggatgag  
 5521 aggggaagtat ccgttcggc ggagatcctg cggaggcca ggaaattccc tcgagcgatg  
 5581 cccatatggg cagcccgga ttacaacct cactgttag agtctggaa ggaccggac  
 5641 tacgtccctc cagtgtgaca cgggtgtcca ttccgcctg ccaaggcccc tcgatacca  
 5701 cctccacgga ggaagaggac ggtgtcctg tcagaatcta ccgtgtctt tgccttggcg  
 20 5761 gagctcgcca caaagacctt cggcagctcc gaatcgtcgg ccgtcgacag cggcacggca  
 5821 acggcctctc ctgaccagcc ctccgacgac ggcgacgagg gatccgacgt tgagctgtac  
 5881 tctccatgc cccccctga gggggagcgg ggggatccc atctcagcga cgggtcttg  
 5941 tctaccgtaa gcgaggaggc tagtgaggac gtcgtctgt gtcgatgtc ctacacatgg  
 6001 acaggcgccc tgatcacgcc atgcgtcgg gaggaacca agtgcctcat caatgcactg  
 25 6061 agcaactctt tgctccgtca ccacaactt gtctatgcta caacatctc cagcgaagc  
 6121 ctgaggcaga agaagggtcac cttgacaga ctgcaggctc tggacgacca ctaccgggac  
 6181 gtgctcaagg agatgaaggc gaaggcgtcc acagttaagg ctaaactct atccgtggag  
 6241 gaagcctgta agctgacgcc cccacattc gccagatcta aatttggcta tggggcaaag  
 6301 gacgtccgga acctatccag caaggccgtt aaccacatcc gctccgtgtg gaaggacttg  
 30 6361 ctggaagaca ctgagacacc aattgacacc accatcatgg caaaaaatga ggtttctgc  
 6421 gtccaaccag agaagggggg ccgcaagcca gtcgcctta tcgtattccc agatttggg  
 6481 gttcgtgtgt gcgagaaaat ggcccttac gatgtgtct ccacctccc tcaggccgtg  
 6541 atgggctctt catacgatt ccaatactct cctggacagc gggtcgagtt cctggtgaat  
 6601 gcctggaaag cgaagaaatg ccctatggg ttgcatatg acaccgctg ttttactca



5 6661 acggctactg agaatgacat ccgtgttgag gagtcaatct accaatgttg tgacttgccc  
 6721 cccgaagcca gacaggccat aaggctcgtc acagagcggc ttacatcgg gggccccctg  
 6781 actaattcta aagggcagaa ctgcggctat cgcgggtgcc gcgcgagcgg tgtactgacg  
 6841 accagctgcg gtaataccct cacatgttac tgaaggccg ctgcggcctg tcgagctgcg  
 6901 aagctccagg actgcacgat gctcgtatgc ggagacgacc ttgtcgttat ctgtgaaagc  
 10 6961 gcgggggaccc aaggaggacga ggcgagccta cgggccttca cggaggctat gactagatac  
 7021 tctgcccccc ctgggggaccc gcccaaacca gaatacgact tggagtgtgat aacatcatgc  
 7081 tctccaatg tgtcagtcgc gcacgatgca tctggcaaaa ggggtgtacta tctaccctg  
 7141 gacccacca ccccccttgc gcgggctgcg tgggagacag ctagacacac tccagtcaat  
 7201 tctgggctag gcaacatcat catgtatgcg ccaccttgt gggcaaggat gatcctgatg  
 15 7261 actcatttct tctcatcct tctagctcag gaacaacttg aaaaagccct agattgtcag  
 7321 atctacgggg cctgttactc cattagacca ctgacctac ctcagatcat tcaacgactc  
 7381 catggcctta gcgcatttct actccatagt tactctccag gtgagatcaa taggggtggct  
 7441 tcatgcctca ggaaacttgg ggtaccgccc ttgcgagtct ggagacatcg ggccagaagt  
 7501 gtccgcgcta ggctactgtc ccaggggggg agggctgcca ctgttgga gtacctcttc  
 20 7561 aactgggcag taaggaccaa gctcaaactc actccaatcc cggctgcgtc ccagtggat  
 7621 ttatccagct ggttcgttgc tggttacagc gggggagaca tatatcacag cctgtctcgt  
 7681 gcccgacccc gctggttcat gtggtgccta ctctacttt ctgtaggggt aggcatttat  
 7741 ctactcccca accgatgaac ggggagctaa aactccagg ccaataggcc atcctgtttt  
 7801 tttccctttt ttttttct tttttttt tttttttt tttctcttt  
 25 7861 ttttctct tttttcct tttttcct tgggtgctcc atcttagccc tagtcacggc  
 7921 tagctgtgaa aggtccgtga gccgcttgac tgcagagagt gctgatactg gcctctctgc  
 7981 agatcaagta ct

### 30 3. HCV Replicon RNA from cell line HCVR8

SEQ ID NO: 3

1 gccagcccc gattgggggc gacactccac catagatcac tcccctgtga ggaactactg  
 61 tcttcacgca gaaagcgtct agccatggcg ttagtatgag tgtcgtgcag cctccaggac  
 121 cccccctccc gggagagcca tagtggtctg cggaaccggt gactacaccg gaattgccag

5 181 gacgaccggg tcctttcttg gatcaaccg ctcaatgcct ggagatttgg gcgtgcccc  
241 gcgagactgc tagccgagta gtgttggtc gcgaaaggcc ttgtgtact gcctgatagg  
301 gtgcttgcca gtgccccggg aggtctcgt gaccgtcac catgagcacg aatcctaaac  
361 ctcaaagaaa aaccaaagg cgcccatga tgaacaaga tggattgcac gcaggttctc  
421 cggccgcttg ggtggagagg ctattcggct atgactgggc acaacagaca atcggctgct  
10 481 ctgatgccgc cgtgttccg cgtcagcgc aggggcgccc ggttctttt gtcaagaccg  
541 acctgtccgg tgccctgaat gaactgcagg acgaggcagc gcggctatcg tggctggcca  
601 cgacgggcgt tccttcgca gctgtgctc acgtgtcac tgaagcggga agggactggc  
661 tgctattggg cgaagtgccg gggcaggatc tcctgtcac tcacctgct cctgccgaga  
721 aagtatccat catggctgat gcaatgcggc ggctgcatac gcttgatccg gctacctgc  
15 781 cattcgacca ccaagcgaac catcgcatc agcgagcacg tactcggatg gaagccggtc  
841 ttgtcatca ggatgatctg gacgaagagc atcaggggct cgcgccagcc gaactgttcg  
901 ccaggctcaa ggcgcgatg cccgacggcg aggatctcgt cgtgacccat ggcgatgcct  
961 gcttgccgaa tatcatggtg gaaaatggc gctttcttg attcatcgac tgtggccggc  
1021 tgggtgtggc ggaccgctat caggacatag cgttggtac cgtgatatt gctgaagagc  
20 1081 ttggcggcga atgggctgac cgcttcctc gtcttacgg tatcgccgt cccgattcgc  
1141 agcgcatcgc ctctatcgc ctcttgacg agttctctg agtttaaca gaccacaacg  
1201 gttccctct agcgggatca attccgccc tctccctcc cccccctaa cgttactggc  
1261 cgaagccgct tggaataagg ccggtgtgcg ttgtctata tgtattttc caccatattg  
1321 ccgtctttt gcaatgtgag ggcccggaaa cctggccctg tctcttgac gagcattcct  
25 1381 aggggtcttt ccctctcgc caaaggaatg caaggtctgt tgaatgtcgt gaaggaagca  
1441 gttcctctgg aagcttctg aagacaaaca acgtctgtag cgacctttg caggcagcgg  
1501 aacccccac ctggcgacag gtgcctctgc ggccaaaagc cacgtgtata agatacacct  
1561 gcaaaggcgg cacaaccca gtgccacgtt gtgagttgga tagttgtgga aagagtcaaa  
1621 tggctctcct caagcgtatt caacaagggg ctgaaggatg cccagaaggt accccattgt  
30 1681 atgggatctg atctggggcc tcggtgcaca tgctttacat gtgtttagc gaggttaaaa  
1741 aacgtctagg cccccgaac cacggggacg tggtttctt ttgaaaaaca cgataatacc  
1801 atggcgccca ttacggccta ctcccaacag acgcgaggcc tacttggctg catcatcact  
1861 agcctcacag gccgggacag gaaccaggtc gagggggagg tccaagtgt ctccaccgca  
1921 acacaatctt tctggcgac ctgcgtcaat ggcgtgtgtt ggactgtcta tcatggtgcc

5 1981 ggctcaaaga ccctgccgg cccaaagggc ccaatcacc aaatgtacac caatgtggac  
 2041 caggacctcg tcggctggca agcgccccc ggggcgcgtt ccttgacacc atgcacctg  
 2101 ggagctcgg acctttactt ggtcacgagg catgccgatg tcattccggt gcgccggcgg  
 2161 ggcgacagca gggggagcct actctcccc agggccgtct cctacttgaa gggctcttcg  
 2221 ggcgggtccac tgctctgcc ctcggggcac gctgtgggca tcttcgggc tgccgtgtgc  
 10 2281 acccgagggg ttggaaggc ggtggacttt gtaccgtcg agtctatggG aaccactatg  
 2341 cgggtcccg tcttcacgga caactcgtcc cctccggccg taccgcagac attccaggtg  
 2401 gccatctac acgccctac tggtagcggc aagagcacta aggtgccggc tgcgtatgca  
 2461 gccaagggt ataagggtgt tgcctgaac ccgtccgtcg ccgccaccct aggttcggg  
 2521 gcgtatatgt ctaaggcaca tggtatcgac ctaacatca gaaccggggg aaggaccatc  
 15 2581 accacgggtg ccccatcac gtactccacc tatggcaagt ttctgccga cgggtggtgc  
 2641 tctgggggcg cctatgacat cataatatgt gatgagtcg actcaactga ctgaccact  
 2701 atcctgggca tcggcacagt cctggaccaa gcggagacgg ctggagcgcg actcgtcgtg  
 2761 ctgccaccg ctacgcctcc gggatcggtc accgtgccac atccaacat cgaggaggtg  
 2821 gctctgtcca gcactggaga aatccccctt tatggcaaag ccaccccat cgagaccatc  
 20 2881 aaggggggga ggcacctcat ttctgccat tccaagaaga aatgtgatga gctgccgcg  
 2941 aagctgtccg gcctcggact caatgtgta gcatattacc ggggccttga tgtatccgtc  
 3001 ataccaacta gcggagacgt cattgtcgt gcaacggacg ctctaagac gggctttacc  
 3061 ggcgatttcg actcagtgat cgactgcaat acatgtgtca ccagacagt cgacttcagc  
 3121 ctggaccga ccttcacat tgagacgacg accgtgccac aagacgggt gtcacgctcg  
 25 3181 cagcggcgag gcaggactgg taggggcagg atgggcattt acaggttgt gactccagga  
 3241 gaacggccct cgggcatgtt cgattcctcg gttctgtcg agtgctatga cgcgggctgt  
 3301 gcttggtacg agctcacgcc cgccgagacc tcagttagggt tgcgggctta cctaaacaca  
 3361 ccagggttgc ccgtctgcca ggaccatctg gattctggg agagcgtctt tacaggcctc  
 3421 acccatag acgccattt cttgtcccag actaagcagg caggagacaa ctccctac  
 30 3481 ctggtagcat accaggctac ggtgtgcgcc agggctcagg ctccacctcc atcgtgggac  
 3541 caaatgtgga agtgtctcat acggctaaag cctacgtgc acgggccaac gccctgtctg  
 3601 tataggctgg gagccgttca aaacgaggtt actaccacac acccataac caaatacatc  
 3661 atggcatgca tgcggctga cctggaggtc gtcacgagca cctgggtgct ggtaggcgga  
 3721 gtcctagcag ctctggccgc gtattgcctg acaacaggca gcgtggtcat tgtgggcagg

5 3781 atcatcttgt ccggaagcc ggccatcatt cccgacaggg aagtccttta ccgggagttc  
 3841 gatgagatgg aagagtgcgc ctcacacctc cttacatcg aacagggaat gcagctcgcc  
 3901 gaacaattca aacagaaggc aatcgggttg ctgcaaacag ccaccaagca agcggaggct  
 3961 gctgctccc tggtggaatc caagtggcgg accctcgaag cttctgggc gaagcatatg  
 4021 tggaattca tcagcgggat acaatattha gcaggctgt cactctgcc tggcaacccc  
 10 4081 gcgatagcat cactgatggc attcacagcc tctatcacca gcccgtcac cacccaacat  
 4141 accctcctgt ttaacatcct ggggggatgg gtggcgccc aacttgctcc tccagcgct  
 4201 gcttctgctt tctagagcgc cggcatcgct ggagcggctg ttggcagcat aggccttggg  
 4261 aagggtcttg tggatatitt ggcaggttat ggagcagggg tggcagcgc gctcgtggcc  
 4321 ttaaaggta tgagcggcga gatgccctcc accgaggacc tggtaacct actccctgct  
 15 4381 atcctctccc ctggcgccct agtcgtcggg gtcgtgtgcg cagcgatact gcgtcggcac  
 4441 gtgggcccag gggagggggc tgtcagtgg atgaaccggc tgatagcgtt cgcttcggg  
 4501 ggtaaccacg tctccccac gcactatgtg cctgagagcg acgtgcagc acgtgtcact  
 4561 cagatctct ctagtctac catcactcag ctgctgaaga ggcttcacca gtggatcaac  
 4621 gaggactgct ccacgccatg ctccggctcg tggctaagag atgtttgga ttggatatgc  
 20 4681 acggtgttga ctgattcaa gacctggctc cagtccaagc tctgccgcg attgccggga  
 4741 gtccccctct tctcatgtca acgtgggtac aaggagctt ggcggggcga cggcatcatg  
 4801 caaaccacct gcccattgg agcacagatc accggacatg tgaaaaaAAAcgg ttccatgagg  
 4864 atcgtggggc ctaggacctg tagtaacacg tggcatggaa cattccccat taacgcgtac  
 4924 accacgggcc cctgcacgcc ctccccggcg ccaattatt ctaggcgct gtggcgggtg  
 25 4984 gctgctgagg agtacgtgga ggttacggg gtgggggatt tccactacgt gacgggcatg  
 5044 accactgaca acgtaaagt cccgtgtcag gttccggccc ccgaattctt cacagaagtg  
 5104 gatggggcgc ggttgacag gtacgtcca gcgtgcaaac cctcctacg ggaggaggtc  
 5164 acattcctgg tggggtcaa tcaatacctg gttgggtcac agctcccatg cgagcccgaa  
 5224 ccggacgtag cagtgtcac ttcatgctc accgaccct cccacattac ggcggagacg  
 30 5284 gctaagcgta ggctggccag gggatctccc cctccttg ccagtcac agtagccag  
 5344 ctgtctgcgc ctctctgaa ggcaaTatgc actaccgtc atgactcccc ggacgtgac  
 5404 ctcatcgagg ccaacctcct gtggcgccag gagatggcg ggaacatcac ccgctggag  
 5464 tcagaaaata aggtagtaat ttggactct ttcagccgc tccaagcga ggaggatgag  
 5524 agggaagtat ccgtccggc ggagatcctg cggaggtcca ggaaattccc tcgagcgtg

5 5584 cccatatggg cacgcccga ttacaacct cactgttag agtcctggaa ggacccggac  
 5644 tacgtccctc cagtgttaca cgggtgtcca ttgccgctg ccaaggcccc tccgatacca  
 5704 cctccacgga ggaagaggac ggtgtcctg tcagaatcta ccgtgtcttc tgccttggcg  
 5764 gagctcgcca caaagacctt cggcagctcc gaatcgtcgg ccgtcgacag cggcacggca  
 5824 acggcctctc ctgaccagcc ctccgacgac ggcgacgagg gatccgacgt tgagtcgtac  
 10 5884 tcctccatgc ccccccttga gggggagccg ggggatccg atctcagcga cgggtcttgg  
 5944 tctaccgtaa gcgaggaggc tagtgaggac gtcgtctgct gctcgtatgc ctacacatgg  
 6004 acaggcggcc tgatcacgcc atgcgctcgg gaggaacca agctgcccat caatgcactg  
 6064 agcaactctt tgcctcgtca ccacaactg gtctatgcta caacatctcg cagcgcaagc  
 6124 ctgcggcgaga agaaggtcac ctttgacaga ctgcaggtcc tggacgacca ctaccgggac  
 15 6184 gtgtcaagg agatgaaggc gaaggcgtcc acagttaagg ctaactct atccgtggag  
 6244 gaagcctgta agctgacgcc cccacattcg gccagatcta aatttggcta tggggcaag  
 6304 gacgtccgga acctatccag caaggccgtt aaccacatcc gctccgtgtg gaaggacttg  
 6364 ctggaagaca ctgagacacc aattgacacc accatcatgg caaaaatga ggttttctgc  
 6424 gtccaaccag agaagggggg ccgcaagcca gtcgcctta tcgtattccc agatttggg  
 20 6484 gttcgtgtgt gcgagaaaat ggccctttac gatgtggtct ccacctccc tcaggccgtg  
 6544 atgggctctt catacggatt ccaatactct cctggacagc gggtcgagtt cctggtgaat  
 6604 gcctggaaag cgaagaaatg ccctatgggc ttgcgatag acaccgctg ttttactca  
 6664 acggtcactg agaatgacat ccgtgttgag gagtcaatct accaatgttg tgacttggcc  
 6724 cccgaagcca gacaggccat aaggctcgtc acagagcggc ttacatcgg gggccccctg  
 25 6784 actaattcta aagggcagaa ctgcggctat cgccgggtgcc gcgcgagcgg tgtactgacg  
 6844 accagctcgg gtaataacct cacatgttac ttgaaggccg ctgcggcctg tcgagctcgg  
 6904 aagctccagg actgcacgat gctcgtatgc ggagacgacc ttgtcgttat ctgtgaaagc  
 6964 gcggggaccc aagaggacga ggcgagccta cgggccttca cggaggctat gactagatac  
 7024 tctgcccccc ctggggaccc gcccaaacca gaatacgact tggagtgtat aacatcatgc  
 30 7084 tcctccaatg tgcagtcgc gcacgatgca tctggcaaaa ggggtgtacta tctacccgt  
 7144 gacccacca ccccccttgc gcgggctcgg tgggagacag ctgacacac tccagtcaat  
 7204 tcctggctag gcaacatcat catgtatgcg cccaccttgt gggcaaggat gatcctgatg  
 7264 actcatttct tctccatct tctagctcag gaacaacttg aaaaagccct agattgtcag  
 7324 atctacgggg cctgttactc cattgagcca cttgacctac ctcatcat tcaacgactc

5      7384 catggcctta ggcgattttc actccatagt tactctccag gtgagatcaa tagggtggct  
       7444 tcatacctca ggaaacttgg ggtaccgccc ttgcgagtct ggagacatcg ggccagaagt  
       7504 gtccgcgcta ggctactgtc ccaggggggg agggctgcca ctgtggcaa gtacctcttc  
       7564 aactgggcag taaggaccaa gctcaaactc actccaatcc cggctgcgtc ccagttggat  
       7624 ttatccagct gggtcgttc tggttacagc gggggagaca tatatcacag cctgtctcgt  
 10      7684 gcccgaaccc gctgggtcat gtggtgccta ctctacttt ctgtaggggt aggcatttat  
       7744 ctactcccca accgatgaac ggggagctaa acactccagg ccaataggcc atcctgtttt  
       7804 ttccctttt tttttctt tttttttt tttttttt tttttttt ttctctttt  
       7864 ttttctct tttttcct ttctttcct tgggtgctcc atcttagccc tagtcacggc  
       7924 tagcttgtaa aggtccgtga gccgcttgac tgcagagagt gctgatactg gcctctctgc  
 15      7984 agatcaagta ct

#### 4. HCV Replicon RNA from cell line HCVR9

SEQ ID NO: 4

20      1 gccagccccc gattgggggc gacactccac catagatcac tcccctgtga ggaactactg  
       61 tcttcacgca gaaagcgtct agccatggcg ttagtatgag tgtcgtgcag cctccaggac  
       121 cccccctccc gggagagcca tagtggctcg cggaaccggg gagtacaccg gaattgccag  
       181 gacgaccggg tcctttcttg gatcaaccgc ctcaatgcct ggagatttgg gcgtgcccc  
       241 gcgagactgc tagccgagta gtgttgggtc gcgaaaggcc ttgtggtact gcctgatagg  
 25      301 gtgcttgcca gtgccccggg aggtctcgta gaccgtgcac catgagcacg aatcctaaac  
       361 ctcaaagaaa aaccaaaggc cgcgccatga ttgaacaaga tggattgcac gcaggttctc  
       421 cggccgcttg ggtggagagg ctattcggct atgactgggc acaacagaca atcggctgct  
       481 ctgatgccgc cgtgttccgg ctgtcagcgc aggggcgccc ggttctttt gtcaagaccg  
       541 acctgtccgg tgccctgaat gaactgcagg acgaggcagc gcggctatcg tggctggcca  
 30      601 cgacgggctg tccttgcgca gctgtgctcg acgtgtcac tgaagcgga agggactggc  
       661 tgctattggg cgaagtgccg gggcaggatc tcctgtcacc tcaccttgc cctgccgaga  
       721 aagtatccat catggctgat gcaatgcggc ggctgcatac gcttgatccg gctacctgcc  
       781 cattcgacca ccaagcgaaa catcgcacg agcgagcacg tactcggatg gaagccggtc  
       841 ttgtcatca ggatgatctg gacgaagagc atcaggggct cgcgccagcc gaactgttcg

5      901 ccaggctcaa ggcgcgcatg cccgacggcg aggatctcgt cgtgacccat ggcgatgcct  
       961 gcttgccgaa tatcatggtg gaaaatggcc gcttttctgg aticatcgac tgtggccggc  
       1021 tgggtgtggc ggaccgctat caggacatag cgttggctac ccgtgatatt gctgaagagc  
       1081 ttggcggcga atgggctgac cgcttctcgt tgccttacgg tatcgccgct cccgattcgc  
       1141 agcgcatcgc ctctatcgc ctcttgacg agttcttctg agtttaaca gaccacaacg  
 10    1201 gtttccctct agcgggatca attcggcccc tctccctccc cccccctaa cgttactggc  
       1261 cgaagccgct tggaataagg ccggtgtgcg ttgtctata tgttatttc caccatattg  
       1321 ccgtctttg gcaatgtgag ggccccgaaa cctggccctg tcttctgac gagcattcct  
       1381 aggggtcttt cccctctcgc caaaggaatg caaggctctg tgaatgtcgt gaaggaagca  
       1441 gttctctgg aagcttctg aagacaaaca acgtctgtag cgacctttg caggcagcgg  
 15    1501 aacccccac ctggcgacag gtgcctcgc ggccaaaagc cacgtgtata agatacacct  
       1561 gcaaaggcgg cacaaccca gtgccagti gtgagtga tagttgtgga aagagtcaaa  
       1621 tggtctcct caagctatt caacaagggg ctgaaggatg cccagaaggt acccattgt  
       1681 atgggatctg atctggggcc tcggtgcaca tgccttcat gtgttagtc gaggttaaaa  
       1741 aacgtctagg cccccgaac cacggggacg tggtttctt ttgaaaaca cgataatacc  
 20    1801 atggcgcta ttacggcta ctcccaacag acgcgaggcc tacttggctg catcatcact  
       1861 agcctcacag gccgggacag gaaccaggtc gagggggagg tccaGgtggt ctccaccga  
       1921 acacaattt tcttggcgac ctgcgtcaat ggcggtgtgt ggactgtcta tcatggtgcc  
       1981 ggctcaaaga ccttgcggg cccaaaggc ccaatcacc aaatgtacac caatgtggac  
       2041 caggacctg tcggctggca agcgcccc gggcgcggt ccttgacacc atgcacctgc  
 25    2101 ggcagctcgg accttactt ggtcacagg catgccgatg tcattccggt gcgccggcgg  
       2161 ggcgacagca gggggagcct actctcccc aggcccgct cctacttga gggtctctg  
       2221 ggcggtcac tgcctgccc ctggggcac gctgtgggca tcttcgggc tgcctgtgc  
       2281 acccgagggg ttgcgaagg ggtggactt gtaccgctg agtctatga aaccactatg  
       2341 cggccccgg tcttcagga caactcgtc cctccggcg taccgagac attccaggtg  
 30    2401 gccatctac acgccctac tggtagcggc aagagcacta aggtgccggc tgcgtatgca  
       2461 gccaagggt ataagtgct tgcctgaac ccgtccgtc ccgccacct aggtttcggg  
       2521 gcgtatatgt ctaaggcaca tggatcgac cctaacaat gaaccgggt aaggaccatc  
       2581 accaggggtg ccccatcac gtactccacc tatggcaagt tcttgccga cgggtgtgc  
       2641 tctggggcg cctatgacat cataatatgt gatgagtgc actcaactga ctcgaccact

5 2701 atcctgggca tcggcacagt cctggaccaa gcggagacgg ctggagcgcg actcgtcgtg  
 2761 ctgccaccg ctacgcctcc gggatcggtc accgtgccac atccaaacat cgaggagggtg  
 2821 gctctgtcca gcaactggaga aatccccctt tatggcaaag ccatccccat cgagaccatc  
 2881 aaggggggga ggcacctcat ttctgccat tccaagaaga aatgtgatga gctcggcgcg  
 2941 aagctgtccg gctcggact caatgctgta gcatattacc ggggccttga tgtatcgtc  
 10 3001 ataccaacta gcggagacgt cattgtcgtg gcaacggacg ctctaagac gggctttacc  
 3061 ggcgatttcg actcagtgat cgactgcaat acatgtgtca cccagacagt cgacttcagc  
 3121 ctggaccga ccttcacat tgagacgacg accgtgccac aagacgcggt gtcacgctcg  
 3181 cagcggcgag gcaggactgg taggggcagg atgggcattt acaggtttgt gactccagga  
 3241 gaacggccct cgggcatgtt cgattcctcg gtctgtgcg agtctatga cgcgggctgt  
 15 3301 gcttggtacg agctcacgcc cgccgagacc tcagttaggt tgcgggctta cctaaacaca  
 3361 ccagggttgc ccgtctgcca ggaccatctg gagtctggg agagcgtctt tacaggcctc  
 3421 accacatag acgcccattt ctgtcccg actaagcagg caggagacaa ctccccctac  
 3481 ctggtagcat accaggctac ggtgtgcgcc agggctcagg ctccacctcc atcgtgggac  
 3541 caaatgtgga agtgtctcat acggctaaag cctacgctgc acgggccaac gccctgctg  
 20 3601 tataggctgg gagccgttca aaacgaggtt actaccacac accccataac caaatacatc  
 3661 atggcatgca tgcggctga cctggaggtc gtcacgagca cctgggtgct ggtaggcgga  
 3721 gtctagcag ctctggccgc gtattgcctg acaacaggca gcgtggtcat tgtgggcagg  
 3781 atcatctgt ccggaaGgcc ggccatcatt cccgacaggg aagtccttta ccgggagttc  
 3841 gatgagatgg aagagtgcgc ctacacctc cttacatcg aacagggaat gcagctcgcc  
 25 3901 gaacaattca aacagaaggc aatcgggttg ctgcaaacag ccaccaagca agcggaggct  
 3961 gctgctcccg tggtggaatc caagtggcgg accctcgaag cttctgggc gaagcatatg  
 4021 tggaaattca tcagcgggat acaatattta gcaggcttgt ccactctgcc tggcaacccc  
 4081 gcgatagcat cactgatggc attcacagcc tctatcacca gcccgctcac cacccaacat  
 4141 accctcctgt ttaacatcct ggggggatgg gtggccgcc aacttgctcc tcccagcgt  
 30 4201 gcttctgctt tcgtaggcgc cgcatcgct ggagcggctg ttggcagcat aggccttggg  
 4261 aaggtgcttg tggataattt ggcaggttat ggagcagggg tggcaggcgc gctcgtggcc  
 4321 ttaaggtca tgagcggcga gatgccctcc accgaggacc tggCtaacct actccctgct  
 4381 atcctctccc ctggcgccct agtcgtcggg gtcgtgtgcg cagcgatact gcgtcggcac  
 4441 gtgggccag gggagggggc tgtcagtgg atgaaccggc tgatagcgtt cgcttcgcgg



5 4501 ggtaaccacg tctccccac gcactatgtg cctgagagcg acgctgcagc acgtgtcact  
 4561 cagatcctct ctagtcttac catcactcag ctgtgaaga ggcttcacca gtggatcaac  
 4621 gaggactgct ccacgccatg ctccggctcg tggctaagag atgtttggga ttggatatgc  
 4681 acgggtgtga ctgattcaa gacctggctc cagtccaagc tcctgccgag attgccggga  
 4741 gtcccttct tctcatgtca acgtgggtac aaggagctt ggcggggcga cggcatcatg  
 10 4801 caaaccacct gcccatgtgg agcacagatc accggacatg tgaaaaaAgg ttccatgagg  
 4861 atcgtggggc ctaggacctg tagtaacacg tggcatggaa cattcccat taacgcgtac  
 4921 accacgggcc cctgcacgcc ctccccggcg ccaattatt ctaggcgct gtggcgggtg  
 4981 gctgctgagg agtacgtgga ggttacgcgg gtgggggatt tccactacgt gacgggcatg  
 5041 accactgaca acgtaaagt cccgtgtcag gttccggccc ccgaattct cacagaagt  
 15 5101 gatggggtgc ggttgacacg gtacgtcca gcgtgcaaac ccctctacg ggaggaggct  
 5161 acattcctgg tcgggtcaa tcaatactg gttgggtcac agtcccatg cgagccgaa  
 5221 ccggacgtag cagtgtcac ttcatgtc accgacctt cccacattac ggcgagacg  
 5281 gtaagcgta gggtggccag gggatctccc ccctcttg ccagtcac agctagccag  
 5341 ctgtctgcgc ctcttgaa ggcaacatgc actaccgtc atgactccc ggacgtgac  
 20 5401 ctcatcgagg ccaacctct gtggcgccag gagatggcg ggaacatcac ccgctggag  
 5461 tcagaaaata aggtagta tttgactct ttcagccgc tccaagcgga ggaggatgag  
 5521 agggaagtat ccgtccgc ggagatcctg cggaggtcca ggaaattccc tcagcgatg  
 5581 cccatattgg cacgcccga ttacaacct cactgttag agtctggaa ggaccggac  
 5641 tacgtccctc cagtgttaca cgggtgtcca ttccgctg ccaagcccc tcgatacca  
 25 5701 cctccacgga ggaagaggac ggtgtcctg tcagaatcta ccgtgtctc tgccttggcg  
 5761 gagctcgcca caaagacct cggcagctc gaatcgtcg ccgtcgacag cggcacggca  
 5821 acggcctctc ctgaccagcc ctccgacgac ggcgacgcgg gatccgacgt tgatcgtac  
 5881 tctccatgc ccccttga gggggagccg ggggatccc atctcagcga cgggtcttg  
 5941 tctaccgtaa gcgaggaggc tagtgaggac gtcgtctgt gtcgatgtc ctacacatg  
 30 6001 acaggcgccc tgatcacgcc atgcgtgag gaggaacca agctgccc ataatgcactg  
 6061 agcaactctt tgcctcgtca ccacaactg gtctatgta caacatctc cagcgcaagc  
 6121 ctgcggcaga agaaggtcac ctgtgacaga ctgcaggtc tggacgacca ctaccgggac  
 6181 gtgctcaagg agatgaaggc gaaggcgtc acagtaagg ctaaactct atccgtggag  
 6241 gaagcctgta agctgacgcc cccacattc gccagatcta aatttggta tggggcaaag

5      6301 gacgtccgga acctatccag caaggccgtt aaccacatcc gtcctgtgtg gaaggacttg  
       6361 ctggaagaca ctgagacacc aattgacacc accatcatgg caaaaatga ggttttctgc  
       6421 gtccaaccag agaagggggg ccgcaagcca gtcgcctta tcgtattccc agatttgggg  
       6481 gtctgtgtgt gcgagaaaat ggccctttac gatgtggtct ccacctccc tcaggccgtg  
       6541 atgggtctct catacggatt ccaatactct cctggacagc gggctgagtt cctggtgaat  
 10     6601 gcctggaaag cgaagaaatg ccctatgggc ttgcatatg acacccgtg ttttactca  
       6661 acggtcactg agaatgacat ccgtgttgag gagtcaatct accaatgttg tgacttgccc  
       6721 cccgaagcca gacaggccat aaggctgctc acagagcggc ttacatcgg gggccccctg  
       6781 actaattcta aagggcagaa ctgctggctat cgccgggtcc gcgcgagcgg tgtactgacg  
       6841 accagctcgc gtaataccct cacatgttac ttgaaggccg ctgctggcctg tcgagctcgc  
 15     6901 aagctccagg actgcacgat gtcgtatgc ggagacgacc ttgtcgttat ctgtgaaagc  
       6961 ggggggaccc aagaggacga ggcgagccta cgggccttca cggaggctat gactagatac  
       7021 tctgcccccc ctggggaccc gcccaaacca gaatacgact tggagtgtat aacatcatgc  
       7081 tctccaatg tgtcagtcgc gcacgatgca tctggcaaaa ggggtgtacta tctcaccctg  
       7141 gacccacca cccccctgc ggggctcgc tgggagacag ctagacacac tccagtcaat  
 20     7201 tcttggttag gcaacatcat catgtatgcg cccacctgt gggcaaggat gatcctgatg  
       7261 actcatttct tctccatcct tctagctcag gaacaacttg aaaaagccct agattgtcag  
       7321 atctacgggg cctgttactc cattgagcca ctgacctac ctcatatcat tcaacgactc  
       7381 catggcctta gcgcatttct actccatagt tactctccag gtgagatcaa taggggtggct  
       7441 tcatgcctca ggaaacttgg ggtaccgccc tgcgagctt ggagacatcg ggccagaagt  
 25     7501 gtccgcgcta ggctactgtc ccaggggggg agggctgcca cttgtggcaa gtacctcttc  
       7561 aactgggcag taaggaccaa gctcaaactc actccaatcc cggctgcgtc ccagttggat  
       7621 ttatccagct ggftcgttgc tggttacagc gggggagaca tatatcacag cctgtctcgt  
       7681 gcccgaaccc gctggttcat gtgtgccta ctctacttt ctgtaggggt aggcattctat  
       7741 ctactcccca accgatgaac ggggagctaa acactccagg ccaataggcc atcctgtttt  
 30     7801 tttcccttt tttttctt tttttttt tttttttt tttctcttt  
       7861 ttttctct ttttctt ttttctt ttttctt ttttctt ttttctt ttttctt  
       7921 tagctgtgaa aggtccgtga gccgcttgac tgcagagagt gctgatactg gcctctctgc  
       7981 agatcaagta ct

5

## 5. HCV Replicon from cell line HCVR22

SEQ ID NO: 5

1 gccagcccc gattgggggc gacactccac catagatcac tccccgtga ggaactactg  
 61 tcttcacgca gaaagcgtct agccatggcg ttagtatgag tgtcgtgcag cctccaggac  
 10 121 cccccctccc gggagagcca tagtgggtcg cggaaccggt gagtacaccg gaattgccag  
 181 gacgaccggg tcctttcttg gatcaaccgc ctcaatgcct ggagatttgg gcgtgcccc  
 241 gcgagactgc tagccgagta gtgtgggtc gcgaaggcc ttgtgtact gcctgatagg  
 301 gtgcttgca gtccccggg aggtctcgt gaccgtgcac catgagcacg aatcctaac  
 361 ctcaaagaaa aaccaaaggc cgcgccatga ttgaacaaga tggattgcac gcaggttctc  
 15 421 cggccgcttg ggtggagagg ctattcggct atgactgggc acaacagaca atcggctgct  
 481 ctgatccgc cgtgtccgc ctgtcagcg aggggcgccc ggttctttt gtcaagaccg  
 541 acctgtccgg tccccgaat gaactgcagg acgaggcagc gcggctatcg tggctggcca  
 601 cgacggcgct tccttgca gctgtgctc acgtgtcac tgaagcggga agggactggc  
 661 tgctattggg cgaagtgcg gggcaggatc tcctgtcatc tcaccttgc cctgccgaga  
 20 721 aagtatccat catggctgat gcaatgcggc ggctgcatac gcttgatccg gctacctgcc  
 781 cattcgacca ccaagcgaat catcgcatcg agcagcacg tactcggatg gaagccggtc  
 841 ttgtcatca ggtatgatc gacgaagagc atcaggggct cgcgccagcc gaactgttcg  
 901 ccaggctcaa ggccgcgatg cccgacggcg aggatctcgt cgtgacctat ggcatgcct  
 961 gcttgccgaa tatcatggtg gaaaatggcc gcttttcttg attcatcgac tgtggccggc  
 25 1021 tgggtgtggc ggaccgctat caggacatag cgttggctac ccgtgatatt gctgaagagc  
 1081 ttggcgccga atgggctgac cgcttcctcg tctttacgg tatcgccgct cccgattcgc  
 1141 agcgcacgc ctctatcgc ctcttgacg agttctctg agtttaaca gaccacaacg  
 1201 gtttcctct agcgggatca attccgccc tctccctcc cccccctaa cgttactggc  
 1261 cgaagccgct tggaataagg ccggtgtgcg ttgtctata tgtattttc caccatattg  
 30 1321 ccgtctttg gcaatgtgag ggcccggaaa cctggccctg tctcttgac gagcattcct  
 1381 aggggtcttt cccctctcg caaaggaatg caaggtctgt tgaatgctg gaaggaagca  
 1441 gttcctctgg aagctcttg aagacaaca acgtctgtg cgaccttg caggcagcgg  
 1501 aacccccac ctggcgacag gtgcctctgc ggccaaaagc cacgtgtata agatacacct  
 1561 gcaaaggcgg cacaaccca gtgccacgt gtgagttgga tagtgtgga aagagtcaa

5 1621 tggctctcct caagcgtatt caacaagggg ctgaaggatg cccagaaggt accccattgt  
 1681 atgggatctg atctggggcc tcggtgcaca tgccttacct gtgtttagtc gaggttaaaa  
 1741 aacgtctagg cccccgaaT cacggggacg tggtttctct tgaanaaaca cgataatacc  
 1801 atgggcctta ttacggccta ctccaacag acgcgaggcc tacttggtg catcatcact  
 1861 agcctcacag gccgggacag gaaccaggtc gagggggagg tccaagtgtt ctccaccgca  
 10 1921 acacaatctt tcctggcgac ctgcgtcaat ggctgtgtt ggactgtcta tcatgtgccc  
 1981 ggctcaaaga cccttgccgg cccaaaggcc ccaatcacc aaatgtacac caatgtggac  
 2041 caggacctcg tcggctggca agcgccccc ggggcgcgtt ccttgacacc atgcacctgc  
 2101 ggcagctcgg accttactt ggtcacgagg catgccgatg tcattccggt gcgccggcgg  
 2161 ggcgacagca gggggagcct actctcccc agggccgtct cctactgaa gggctcttcg  
 15 2221 ggcgtccac tgctctgcc ctggggcac gctgtgggca tcttcgggc tcgctgtgc  
 2281 acccgagggg ttgcgaaggc ggtggacttt gtaccgctg agtctatggG aaccactatg  
 2341 cggctcccgg tcttcacgga caactcgtcc cctccggccg taccgcagac attccagggt  
 2401 gccatctac acgcccctac tggtagcggc aagagcacta aggtgccggc tgcgtatgca  
 2461 gccaagggt ataagggtct tgcctgaac ccgtccgtc cgccaccct aggttcggg  
 20 2521 gcgtatatgt ctaaggcaca tggatcgac cctaactca gaaccggggt aaggaccatc  
 2581 accacgggtg ccccatcac gtactccacc tatggcaagt ttctgccga cgggtggtgc  
 2641 tctgggggcg cctatgacat cataatatgt gatgagtgc actcaactga ctgcaccat  
 2701 atctgggca tcggcacagt cctggacaa gcggagacgg ctggagcgc actcgtcgtg  
 2761 ctgccaccg ctacgctcc gggatcggt accgtgccac atccaaacat cgaggagggt  
 25 2821 gctctgtcca gactggaga aatccccctt tatggcaaag ccatcccat cgagaccatc  
 2881 aaggggggga ggcacctcat ttctgcat tccaagaaga aatgtgatga gctcggcgcg  
 2941 aagctgtccg gcctcggact caatgctga gcatattacc ggggccttga tcatccgtc  
 3001 ataccaacta gcggagacgt catgtcgt gcaacggacg ctctaatac gggctttacc  
 3061 ggcgattcg actcagtgt cgactgcaat acatgtgtca cccagacagt cgacttcagc  
 30 3121 ctggaccga ccttcacat tgagacgac accgtgccac aagacgcggt gtcacgctcg  
 3181 cagcggcgag gcaggactgg taggggcagg atgggcattt acaggtttgt gactccagga  
 3241 gaacggccct cgggcatgtt cgattcctg gtctgtgcg agtgctatga cgcgggctgt  
 3301 gcttggtacg agctcacgc cgcgagacc tcagttaggt tgcgggctta cctaacaca  
 3361 ccagggttc ccgtctgcca ggacctctg gatttctggg agagcgtctt tacaggcctc

5 3421 acccacatag acgcccattt ctgtcccag actaagcagg caggagacaa ctcccctac  
 3481 ctggtagcat accaggctac ggtgtgcgc agggctcagg ctccacctcc atcgtgggac  
 3541 caaatgtgga agtgtctcat acggctaaag cctacgtgc acgggccaac gcccctgctg  
 3601 tataggctgg gagccgtica aaacgaggtt actaccacac accccataac caatacatc  
 3661 atggcatgca tgtcggctga cctggaggtc gtcacgagca cctgggtgct ggtaggcgga  
 10 3721 gtcctagcag ctctggccgc gtattgcctg acaacaggca gcgtggtcat tgtgggcagg  
 3781 atcatcttgt ccggaaagcc ggccatcatt cccgacaggg aagtccttta ccgggagttc  
 3841 gatgagatgg aagagtgcgc ctcacacctc cttacatcg aacagggaat gcagctcgcc  
 3901 gaacaattca aacagaaggc aatcgggttg ctgcGaacag ccaccaagca agcggaggct  
 3961 gctgctcccg tgggtgaatc caagtggcgg accctcgaag cttctgggc gaagcatatg  
 15 4021 tggaattica tcagcgggat acaatattha gcaggttgt cactctgcc tggcaacccc  
 4081 gcgatagcat cactgatggc attcacagcc tctatacca gcccgctcac cacccaacat  
 4141 accctctgt ttaacatcct ggggggatgg gtggccgcc aacttgctcc tccagcgct  
 4201 gcttctgctt tcgtaggcgc cggcatcgct ggagcggctg ttggcagcat aggccttggg  
 4261 aaggtgcttg tggatathtt ggcaggttat ggagcagggg tggcagcgcc gctcgtggcc  
 20 4321 ttaaggtca tgagcggcga gatgccctcc accgaggacc tggtaacct actccctgct  
 4381 atcctctccc ctggcgccct agtcgtcggg gtcgtgtcg cagcgatact gcgtcggcac  
 4441 gtgggcccag gggagggggc tgtgcagtgg atgaaccggc tgatagcgtt cgcttcgagg  
 4501 ggtaaccacg tctccccac gcactatgtg cctgagagcg acgctgcagc acgtgtcact  
 4561 cagatcctct ctagtcttac catcactcag ctgctgaaga ggcttcacca gtggatcaac  
 25 4621 gaggactgct ccacgccatg ctccggctcg tggctaagag atgtttggga ttggatatgc  
 4681 acggtgttga ctgattcaa gacctggctc cagtccaagc tctgcccgcg attgccggga  
 4741 gtccccttct tctcatgtca acgtgggtac aaggaggtct ggcggggcga cggcatcatg  
 4801 caaaccacct gcccatgtgg agcacagatc accggacatg tgaaaaacgg ttccatgagg  
 4861 atcgtggggc ctaggacctg tagtaacacg tggcatggaa cattccccat taacgcgtac  
 30 4921 accacgggcc cctgcacgcc ctccccggcg ccaaattatt ctaggcgct gtggcgggtg  
 4981 gctgctgagg agtacgtgga ggttacgcgg gtgggggatt tccactacgt gacgggcatg  
 5041 accactgaca acgtaaagtg cccgtgtcag gttccggccc ccgaattctt cacagaagtg  
 5101 gatggggtgc ggttgacacg gtacgtcca gcgtgcaaac cctcctacg ggaggaggtc  
 5161 acattcctgg tcgggtcaa tcaatactg gttgggtcac agctcccatg cgagcccga

5 5221 ccggacgtag cagtgtctac ttccatgctc accgaccctt cccacattac ggccggagacg  
 5281 gctaagcgta ggctggccag gggatctccc cctccttg ccagctcatc agtagccag  
 5341 ctgtctgcgc ctctcttgaa ggcaacatgc actaccgtc atgactcccc ggacgtgac  
 5401 ctatcgagg ccaacctctt gtggcggcag gagatggcg ggaacatcac ccgctggag  
 5461 tcagaaaata aggtagtaat ttggactct ttgagccgc tccaagcgga ggaggatgag  
 10 5521 aggggaagtat ccgtccggc ggagatcctg cggaggtcca ggaattccc ttgagcgatg  
 5581 cccatatggg cagccccgga ttacaacctt ccaactgttag agtctggaa ggaccggac  
 5641 tacgtccctc cagtgttaca cgggtgtcca ttgccgctg ccaaggcccc tccgatacca  
 5701 cctccacgga ggaagaggac ggttgtctg tcagaatcta ccgtgtctt tgccttggcg  
 5761 gagctgccca caaagacctt cggcagctcc gaatcgctgg ccgtcgacag cggcacggca  
 15 5821 acggcctctc ctgaccagcc ctccgacgac ggcgacgcgg gatccgacgt tgagtctac  
 5881 tctccatgc cccccctga gggggagccg ggggatccg atctcagca cgggtcttg  
 5941 tctaccgtaa gcgaggagc tagtgaggac gtctgtgtg gctgatgtc ctacacatgg  
 6001 acaggcggcc tgatcacgcc atgcgctgcg gaggaacca agtgcctat caatgcactg  
 6061 agcaactctt tgctccgtca ccacaacttg gtctatgcta caacatctg cagcgcaagc  
 20 6121 ctgcggcaga agaaggtcac ctctgacaga ctgcaggtcc tggacgacca ctaccgggac  
 6181 gtgtcaagg agatgaaggc gaaggcgtcc acagttaagg cttaaactct atccgtggag  
 6241 gaagcctgta agctgacgcc cccacattcg gccagatcta aatttggcta tggggcaaag  
 6301 gacgtccgga acctatccag caaggccgtt aaccacatcc gctccgtgtg gaaggacttg  
 6361 ctggaagaca ctgagacacc aattgacacc accatcatgg caaaaaatga ggttttctgc  
 25 6421 gtccaaccag agaagggggg ccgcaagcca gctcgcccta tctattccc agatttggg  
 6481 gttcgtgtg gcgagaaaat ggccctttac gatgtgtct ccacctccc tcaggccgtg  
 6541 atgggctctt cctacggatt ccaatactct cctggacagc ggtcgagtt cctggtgaat  
 6601 gcctggaaag cgaagaaatg ccctatgggc ttgcatatg acaccgctg ttttactca  
 6661 acggtcactg agaatgacat ccgtgttgag gactcaatct accaatgtt tgacttggcc  
 30 6721 cccgaagcca gacaggccat aaggtcgctc acagagcggc ttacatcgg gggccccctg  
 6781 actaattcta aaggcgagaa ctgcggctat cggcggtgcc gcgagagcg tgtactgacg  
 6841 accagctgcg gtaataacct cacatgttac ttgaaggccg ctgcggcctg tcgagctgcg  
 6901 aagctccagg actgcacgat gctcgtatgc ggagacgacc ttgtcgttat ctgtgaaagc  
 6961 gcggggaccc aagaggacga ggcgagccta cgggccttca cggaggctat gactagatac

5 7021 tctgcccccc ctggggaccc gcccaaacca gaatacgact tggagttgat aacatcatgc  
 7081 tctccaatg tgcagtcgc gcacgatgca tctggcaaaa ggggtgtacta tctacccgt  
 7141 gacccacca ccccccttgc gcgggctgcg tgggagacag ctacacacac tccagtcaat  
 7201 tcttggttag gcaacatcat catgtatgcg cccacctgtg gggcaaggat gatcctgatg  
 7261 actcatttct tctccatcct tctagctcag gaacaacttg aaaaagccct agattgtcag  
 10 7321 atctacgggg cctgttactc cattgagcca ctgacctac ctacatcat tcaacgactc  
 7381 catggcctta gcgcatttc actccatagt tactctccag gtgagatcaa tagggtggct  
 7441 tcatgcctca ggaaacttgg ggtaccgccc ttgcgagtct ggagacatcg ggccagaagt  
 7501 gtccgcgcta ggctactgtc ccaggggggg agggctgcca ctgttgcaa gtacctcttc  
 7561 aactgggcag taaggaccaa gctcaaactc actccaatcc cggctgcgct ccagttggat  
 15 7621 ttatccagct ggttcgttgc tggttacagc gggggagaca tatatcacag cctgtctcgt  
 7681 gcccagcccc gctggttcat gtggtgccta ctctacttt ctgtaggggt aggcattctat  
 7741 ctactcccca accgatgaac ggggagctaa aactccagg ccaataggcc atcctgtttt  
 7801 tttccctttt tttttctt tttttttt tttttttt tttctcttt  
 7861 ttttctct tttttctt ttcttctt tgggtgctcc atcttagccc tagtcacggc  
 20 7921 tagctgtgaa aggtccgtga gccgcttgac tgcagagagt gctgatactg gcctctctgc  
 7981 agatcaagta ct

# 6. HCV Replicon from cell line HCVR24

SEQ ID NO: 6

25 1 gccagcccc gattgggggc gacactccac catagatcac tcccctgtga ggaactactg  
 61 tcttcacgca gaaagcgtct agccatggcg ttagtatgag tgtcgtgcag cctccaggac  
 121 cccccctccc gggagagcca tagtggctcg cggaaccggg gagtacaccg gaattgccag  
 181 gacgaccggg tctttcttg gatcaaccog ctcaatgcct ggagatttgg gcgtgcccc  
 241 gcgagactgc tagccgagta gtgttgggtc gcgaaaggcc ttgtgtact gcctgatagg  
 30 301 gtgcttgcga gtgccccggg aggtctcgtg gaccgtgcac catgagcacg aatcctaacc  
 361 ctcaaagaaa aaccaaaggc cgcgccatga ttgaacaaga tggattgcac gcaggttctc  
 421 cggccgcttg ggtggagagg ctattcggct atgactgggc acaacagaca atcggctgct  
 481 ctgatccgc cgtgttcgg ctgtcagcg aggggcgccc ggttctttt gtcaagaccg  
 541 acctgtccgg tgcctgaat gaactgcagg acgaggcagc gcggctatcg tggctggcca

5      601 cgacgggcgt tccttgcgca gctgtgctcg acgttgtcac tgaagcggga agggactggc  
       661 tgctattggg cgaagtgccg gggcaggatc tcctgtcatc tcacctgtc cctgccgaga  
       721 aagtatccat catggctgat gcaatgcggc ggctgcatac gcttgatccg gctacctgcc  
       781 cattcgacca ccaagcgaac catcgcatcg agcagacacg tactcggatg gaagccggtc  
       841 ttgtcgatca ggaatgctg gacgaagagc atcaggggct cgcgccagcc gaactgttcg  
 10     901 ccaggctcaa ggcgcgcatg cccgacggcg aggatctcgt cgtgacctat ggcatgctc  
       961 gcttgccgaa tatcatgggt gaaaatggcc gcttttctgg attcatcgac tgtggccggc  
      1021 tgggtgtggc ggaccgctat caggacatag cgttggctac ccgtgatatt gctgaagagc  
      1081 ttggcggcga atgggctgac cgcttctcgt tgctttacgg tatcgccgct cccgattcgc  
      1141 agcgcacatc ctctatcgc ctcttgacg agttctctg agtttaaaca gaccacaacg  
 15     1201 gtttccctct agcgggatca attccgcccc tctccctccc cccccctaa cgttactggc  
      1261 cgaagccgct tggaataagg ccggtgtgctg ttgtctata tgtattttc caccatattg  
      1321 ccgtctttg gcaatgtgag ggcccggaaa cctggccctg tcttctgac gagcattcct  
      1381 aggggtcttt cccctctcgc caaaggaatg caaggtctgt tgaatgtcgt gaaggaagca  
      1441 gtctctctgg aagcttctg aagacaaaca acgtctgtg cgacctttg caggcagcgg  
 20     1501 aacccccac ctggcgacag gtgcctctgc ggccaaaagc cacgtgtata agatacacct  
      1561 gcaaaggcgg cacaaccca gtgccacgtt gtgagtggga tagttgtgga aagagtcaaa  
      1621 tggctctcct caagcgtatt caacaagggg ctgaaggatg cccagaaggt acccattgt  
      1681 atgggatctg atctggggcc tcggtgcaca tgccttacct gtgtttagtc gaggttaaaa  
      1741 aacgtctagg cccccgaac cagggggacg tggtttctct ttgaaaaaca cgataatacc  
 25     1801 atggcgccca ttacggccta ctcccaacag acgcgaggcc tacttgctg catcatcact  
      1861 agcctcacag gccgggacag gaaccaggtc gagggggagg tccaagtgt ctccaccgca  
      1921 acacaatctt tcctggcgac ctgcgtcaat ggctgtgtt ggactgtcta tcatggtgcc  
      1981 ggctcaaaga cccttgccgg cccaaagggc ccaatcacc aaatgtacac caatgtggac  
      2041 caggacctcg tcggctggca agcggccccc ggggcgcgtt ccttgacacc atgcacctgc  
 30     2101 ggcagctcgg accttactt ggtcacgagg catccgatg tcattccggt gcgccggcgg  
      2161 ggcgacagca gggggagcct actctcccc agggccgtct cctactgaa gggctcttcg  
      2221 ggcggtccac tgctctgccc ctgggggac gctgtgggca tcttcgggc tgccgtgtgc  
      2281 acccgagggg ttgcgaaggc ggtggacttt gtaccgctg agtctatgga aacctatg  
      2341 cggccccggg tcttcacgga caactgtcc cctccggccg taccgcagac attccaggtg



5     2401 gcccatctac acgcccctac tggtagcggc aagagcacta aggtgccggc tgcgtatgca  
       2461 gccaagggt ataagggtct tgcctgaac ccgtccgtcg ccgccaccct aggtttcggg  
       2521 gcgtatatgt ctaaggcaca tggtagcgac cctaactca gaaccggggt aaggaccatc  
       2581 accacgggtg cccccatcac gtactccacc tatggcaagt ttctgccga cgggtggtgc  
       2641 tctgggggcg cctatgacat cataatatgt gatgagtgcc actcaactga ctgaccact  
 10    2701 atcctgggca tcggcacagt cctggaccaa gcggagacgg ctggagcgcg actcgtcgtg  
       2761 ctgccaccg ctacgcctcc gggatcggtc accgtgccac atccaaacat cgaggaggtg  
       2821 gctctgtcca gactggaga aatccccctt tatggcaaag ccatccccat cgagaccatc  
       2881 aaggggggga ggcacctcat ttctgccat tccaagaaga aatgtgatga gctcgcgcg  
       2941 aagctgtccg gcctcgact caatgctgta gcatattacc ggggccttga tgtatccgtc  
 15    3001 ataccaacta gcggagacgt cattgtcgta gcaacggacg ctctaatgac gggctttacc  
       3061 ggcgatttcg actcagtgat cgactgcaat acatgtgtca ccagacagc cgacttcagc  
       3121 ctggaccgga ccttcacat tgagacgacg accgtgccac aagacgcggt gtcacgctcg  
       3181 cagcggcgag gcaggactgg taggggcagg atgggcattt acaggtttgt gactccagga  
       3241 gaacggccct cgggcatgtt cgattcctcg gttctgtcg agtgctatga cgcgggctgt  
 20    3301 gcttggtagc agctcacgcc cgccgagacc tcagttaggt tgcgggctta cctaaacaca  
       3361 ccagggttgc ccgtctgcca ggaccatctg gagttctggg agagcgtctt tacaggcctc  
       3421 acccatag acgcccattt ctgtcccag actaagcagg caggagacaa ctcccctac  
       3481 ctggtagcat accaggctac ggtgtgcgcc agggctcagg ctccacctcc atcgtgggac  
       3541 caaatgtgga agtgtctcat acggctaaag cctacgtgc acgggccaac gccctgctg  
 25    3601 tataggctgg gagccgttca aaacgaggtt actaccacac accccataac caatacatc  
       3661 atggcatgca tgcgggtga cctggaggtc gtcacgagca cctgggtgct gtaggcgga  
       3721 gtcctagcag ctctggccgc gtattgcctg acaacaggca gcgtggtcat tgtgggcagg  
       3781 atcatcttgt ccggaaagcc ggccatcatt ccgacaggg aagtccttta ccgggagttc  
       3841 gatgagatgg aagagtgcgc ctacacctc cttacatcg aacagggaat gcagctcgcc  
 30    3901 gaacaattca aacagaaggc aatcgggttg ctgcaaacag ccaccaagca agcggaggct  
       3961 gctgctcccg tgggtgaatc caagtggcg accctcgaag cttctgggc gaagcatatg  
       4021 tggaatttca tcagcgggat acaatattha gcaggcttgt ccactctgcc tggcaacccc  
       4081 gcgatagcat cactgatggc attcacagcc tctatacca gcccgtcac cacccaacat  
       4141 accctctgt ttaacatcct ggggggatgg gtggcgccc aacttgctcc tccagcgct

5 4201 gcttctgctt tcgtaggcgc cggcatcgct ggagcggctg ttggcagcat aggccttggg  
 4261 aaggtgcttg tggatattt ggcaaggttat ggagcagggg ttggcagggc gctcgtggcc  
 4321 ttaaggta tgagcggcga gatgccctcc accgaggacc tggtaacct actccctgct  
 4381 atcctctccc ctggcgccct agtcgtcggg gtcgtgtgcg cagcgatact ggcgcggcac  
 4441 gtgggcccag gggagggggc tgtgcagtgg atgaaccggc tgatagcgti cgcttcgcg  
 10 4501 ggtaaccacg tctcccccac gcactatgtg cctgagagcg acgtgcagc acgtgtcact  
 4561 cagatcctct ctagtcttac catcactcag ctgctgaaga ggcttcacca gtggatcaac  
 4621 gaggactgct ccacgccatg ctccggctcg tggctaagag atgtttggga ttggatatgc  
 4681 acggtgttga ctgattcaa gacctggctc cagtccaagc tctgcccgcg attgccggga  
 4741 gtccctctct tctcatgtca acgtgggtac aaggaggtct ggccggggcga cggcatcatg  
 15 4801 caaaccacct gcccatgtgg agcacagatc accggacatg tgaaaaacgg ttccatgagg  
 4861 atcgtggggc ctaggacctg tagtaacacg tggcatggaa cattcccat taacgcgtac  
 4921 accacggggc cctgcacgcc ctccccggcg ccaattatt ctaggcgct gtggcgggtg  
 4981 gctgctgagg agtacgtgga ggttacggcg gtgggggatt tccactacgt gacgggcatg  
 5041 accactgaca acgtaaagt cccgtgtcag gtccggccc ccgaattctt cacagaagtg  
 20 5101 gatgggggtg ggttgacacg gtacgtcca gcgtgcaaac cctcctacg ggaggaggtc  
 5161 acattctctg tcgggctcaa tcaatacctg gttgggtcac agctcccatg cgagccgaa  
 5221 ccggacgtag cagtgtcac ttccatgtc accgacctt cccacattac ggcggagacg  
 5281 gctaagcgta ggctggccag gggatctccc cctccttgg ccagtcac agctaTccag  
 5341 ctgtctgcgc ctctctgaa ggcaacatgc actaccgtc atgactccc ggacgtgac  
 25 5401 ctcatcgagg ccaacctct gtggcggcag gagatggcg ggaacatcac ccgctggag  
 5461 tcagaaaata aggtagtaat ttggactct ttgagccgc tccaagcgga ggaggatgag  
 5521 agggaagtat ccgtccggc ggagatcctg cggaggtcca ggaaattccc tcgagcgatg  
 5581 cccatattgg cagccccga ttacaacct cactgttag agtctggaa ggaccggac  
 5641 tacgtccctc cagtgttaca cgggtgtcca ttccgcctg ccaaggcccc tccgatacca  
 30 5701 cctccacgga ggaagaggac ggtgtcctg tcagaatcta ccgtgtctt tgccttggcg  
 5761 gagctcgcca caaagacct cggcagctcc gaatcgtcg ccgtcgacag cggcacggca  
 5821 acggcctctc ctgaccagcc ctccgacgac ggcgacggg gatccgacgt tgagtcgtac  
 5881 tctccatgc cccccctga gggggagccg ggggatccc atctcagcga cgggtcttgg  
 5941 tctaccgtaa gcgaggaggc tagtgaggac gtcgtctgct gctcgatgc ctacacatgg

5      6001 acaggcgccc tgatcacgcc atgcgctgcg gaggaaacca agctgccc ataatgcactg  
       6061 agcaactctt tgctccgtca ccacaacttg gtctatgcta caacatctcg cagcgcaagc  
       6121 ctgcggcaga agaagggtac ctttgacaga ctgcagggtcc tggacgacca ctaccgggac  
       6181 gtgctcaagg agatgaaggc gaaggcgtcc acagttaagg ctaaacttct atccgtggag  
       6241 gaagcctgta agctgacgcc cccacattcg gccagatcta aatttggtta tggggcaaa  
 10     6301 gacgtccgga acctatccag caaggccgtt aaccacatcc gctccgtgtg gaaggacttg  
       6361 ctggaagaca ctgagacacc aattgacacc accatcatgg caaaaaatga ggttttctgc  
       6421 gtccaaccag agaagggggg ccgaagcca gtcgcctta tctattccc agatttggg  
       6481 gtctgtgtgt gcgagaaaat ggccctttac gatgtgtctt ccacctccc tcaggccgtg  
       6541 atgggctctt catacggatt ccaatactct cctggacagc gggtcgagtt cctggtgaat  
 15     6601 gcctggaaag cgaagaaatg ccctatgggc ttgcatatg acaccgctg ttttactca  
       6661 acggtcactg agaatgacat ccgtgttgag gactcaatct accaatgttg tgacttggcc  
       6721 cccgaagcca gacaggccat aaggtcgctc acagagcggc ttacatcgg gggcccccctg  
       6781 actaattcta aagggcagaa ctgcggctat cgccgggtgc gcgcgagcgg tgtactgacg  
       6841 accagctgcg gtaataccct cacatgttac tgaaggccg ctgcggcctg tcgagctgcg  
 20     6901 aagctccagg actgcacgat gctcgtatgc ggagacgacc ttgtcgttat ctgtgaaagc  
       6961 gcggggaccc aagaggacga ggcgagccta cgggccttca cggaggctat gactagatac  
       7021 tctgcccccc ctggggaccc gccc aaacca gaatacgact tggagtgtat aacatcatgc  
       7081 tctccaatg tctcagtcgc gcacgatgca tctggcaaaa ggtgttacta tctcaccgt  
       7141 gacccacca ccccccttc gcgggctgcg tgggagacag ctgacacac tccagtcaat  
 25     7201 tcttggttag gcaacatcat catgtatgcg cccacctgtt gggcaaggat gatcctgatg  
       7261 actcatttct tctccatct tctagctcag gaacaacttg aaaaagccct agattgtcag  
       7321 atctacgggg cctgttactc cattgagcca cttgacctac ctcagatcat tcaacgactc  
       7381 catggcctta gcgcatttct actccatagt tactctccag gtgagatcaa tagggtggct  
       7441 tcatgcctca ggaaacttgg ggtaccgccc ttgcgagtct ggagacatcg ggccagaagt  
 30     7501 gtccgcgcta ggctactgtc ccaggggggg agggctgcca cttgtggcaa gtaccttctc  
       7561 aactgggcag taaggaccaa gctcaaaact actccaatcc cggctgcgtc ccagtggat  
       7621 ttatccagct ggttcgttgc tggttacagc gggggagaca tatatcacag cctgtctcgt  
       7681 gcccgacccc gctggttcat gtggtgccta ctctacttt ctgtagggtt aggcattctat  
       7741 ctactcccca accgatgaac ggggagctaa aactccagg ccaataggcc atcctgtttt

5     7801 ttccctttt ttttttctt tttttttt tttttttt tttttttt ttctctttt  
      7861 tttttctct ttttttctt ttctttctt tgggggtcc atcttagccc tagtcacggc  
      7921 tagctgtgaa aggtccgtga gccgcttgac tgcagagagt gctgatactg gcctctctgc  
      7981 agatcaagta ct

SEQ ID NO: 7

10    GCCCTAGATTGTCAGATCTACG

SEQ ID NO: 8

ATAAATCCAACTGGGACGCAGC

15    SEQ ID NO: 9

CCTTGTGGGCAAGGATGATCC

SEQ ID NO: 10

GACAGGCTGTGATATATGTCTCC

20

SEQ ID NO: 11

GTTCTTTTTGTCAAGACCGACC

SEQ ID NO: 12

25    CCACCATGATATTCGGCAAGC

SEQ ID NO: 13

CTCGTATGTTGTGTGGAA

30    SEQ ID NO: 14

GTCGCTCTCGAGGCACATA

## WHAT IS CLAIMED IS:

- 1 1. An isolated nucleic acid molecule encoding a replication competent recombinant  
2 Hepatitis C Virus (HCV) genome, which nucleic acid comprises all or part of an HCV  
3 genome and is able to replicate efficiently when transfected into a susceptible cell line  
4 without reducing the growth rate of said cell line by more than 10 fold.
- 1 2. The isolated nucleic acid molecule encoding a recombinant HCV genome of claim 1,  
2 which nucleic acid comprises from 5' to 3' on the positive-sense nucleic acid  
3 (a) a functional 5' HCV non-translated region (NTR) comprising an extreme  
4 5'-terminal conserved sequence;  
5 (b) at least one open reading frame (ORF) encoding a heterologous gene  
6 operatively associated with an expression control sequence, wherein the  
7 heterologous gene and expression control sequence are oriented on the  
8 positive-strand nucleic acid molecule;  
9 (c) an ORF encoding at least a portion of an HCV polypeptide whose cleavage  
10 products form functional components of HCV virus particles and RNA  
11 replication machinery, and  
12 (d) an HCV 3' NTR comprising an extreme 3'-terminal conserved sequence,  
13 and wherein said nucleic acid is able to replicate efficiently in a susceptible  
14 cell line without reducing the growth rate of said cell line by more than 10  
15 fold.
- 1 3. The isolated nucleic acid of claim 1, wherein the susceptible cell line is selected from  
2 the group consisting of human hepatoma cell line Huh-7, human hepatoma cell line  
3 HepG2, hepatoma cell line PH5CH, *T. belangeri* liver cell line MBTL, human diploid  
4 fibroblast cell line VERO, secondary monkey kidney cell line CV-1, T cell line MT-2,  
5 T cell line HPBMa10-2, T cell line MOLT-4, and B cell line Daudi.

- 1 4. The susceptible cell line of claim 4, which is human hepatoma cell line Huh-7.
- 1 5. The isolated nucleic acid molecule according to claim 1, which is selected from the  
2 group consisting of double stranded DNA, single stranded DNA, double stranded  
3 RNA, and single stranded RNA.
- 1 6. An isolated nucleic acid molecule which is not more than 99.9% identical and is at  
2 least 95% identical to SEQ ID NO: 1.
- 1 7. The isolated nucleic acid molecule of claim 6 comprising nucleotide sequence of  
2 HCVR 2 (SEQ ID NO: 2).
- 1 8. The isolated nucleic acid molecule of claim 6 comprising nucleotide sequence of  
2 HCVR 8 (SEQ ID NO: 3).
- 1 9. The isolated nucleic acid molecule of claim 6 comprising nucleotide sequence of  
2 HCVR 9 (SEQ ID NO: 4).
- 1 10. The isolated nucleic acid molecule of claim 6 comprising nucleotide sequence of  
2 HCVR 22 (SEQ ID NO: 5).
- 1 11. The isolated nucleic acid molecule of claim 6 comprising nucleotide sequence of  
2 HCVR 24 (SEQ ID NO: 6).
- 1 12. A stable cell line transfected with the isolated nucleic acid molecule according to  
2 claim 1, wherein said cell line:

- 3 (a) has a growth rate which is not less than 10% of the growth rate of the  
4 corresponding naïve cell line, and  
5 (b) is capable of supporting efficient replication of said isolated nucleic acid.

- 1 13. The cell line of claim 12 wherein said cell line is selected from the group consisting of  
2 human hepatoma cell line Huh-7, human hepatoma cell line HepG2, hepatoma cell  
3 line PH5CH, *T. belangeri* liver cell line MBTL, human diploid fibroblast cell line  
4 VERO, secondary monkey kidney cell line CV-1, T cell line MT-2, T cell line  
5 HPBMa10-2, T cell line MOLT-4, and B cell line Daudi.
- 1 14. The cell line of claim 12 wherein said cell line is derived from a human hepatoma cell  
2 line Huh-7.
- 1 15. The cell line of claim 14 designated HCVR 2 and having ATCC Accession No. PTA-  
2 2489.
- 1 16. The cell line of claim 14 designated HCVR 8 and having ATCC Accession No. PTA-  
2 2490.
- 1 17. The cell line of claim 14 designated HCVR 9 and having ATCC Accession No. PTA-  
2 2486.
- 1 18. The cell line of claim 14 designated HCVR 22 and having ATCC Accession No.  
2 PTA-2487.
- 1 19. The cell line of claim 14 designated HCVR 24 and having ATCC Accession No.  
2 PTA-2488.


- 1 20. A method of screening for anti-HCV therapeutics, which method comprises  
2 comparing a level of HCV subgenomic replicon RNA or replicon RNA-associated  
3 protein expression in the cell line of claim 12 contacted with a candidate therapeutic  
4 agent to the cell line not contacted with the candidate therapeutic agent, wherein a  
5 decrease in the level of HCV subgenomic replicon RNA or replicon RNA-associated  
6 protein expression is indicative of the inhibitory activity of the agent.
- 1 21. A method for detecting antibodies to HCV in a biological sample from a subject  
2 comprising contacting said sample with the protein fractions derived from the cell line  
3 of claim 12 under conditions that permit interaction of HCV-specific antibodies in the  
4 sample with the HCV protein(s) produced in said cell line, followed by detecting  
5 binding of the antibodies in the sample to these HCV-derived protein(s), wherein said  
6 binding is indicative of the presence of HCV infection in the subject from which the  
7 sample was derived.
- 1 22. The method of claim 21 wherein said biological sample is selected from the group  
2 consisting of blood, serum, plasma, blood cells, lymphocytes, and liver cells.






**FIG. 1**

PRIOR ART



1/4

**FIG. 3**



**FIG. 2**

PRIOR ART




FIG. 4





FIG. 5



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US01/46350

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :C12Q 1/68; G01N 33/53; C12N 5/00; C07H 21/04  
US CL :435/6, 7.1, 325; 536/23.2

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 7.1, 325; 536/23.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

DIALOG, MEDLINE, APS, HCV, replication competent recombinant, growth rate, cell line, Huh-7, therapeutics

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,874,565A (RICE et al) 23 February 1999, claims.	1-5, 12-14, 20-22

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

"	Special categories of cited documents:	"I"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E"	earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Z"	document member of the same patent family
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

25 MARCH 2002

Date of mailing of the international search report

01 MAY 2002

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

*Valerie Bell-Harris for*  
HANKYEL TIMOTHY PARK, PH.D.

Telephone No. (703) 308-0196